arXiv:0907.5305v1 [math.PR] 30 Jul 2009

A historical law of large numbers for the Marcus-Lushnikov process

Stéphanie Jacquot[*]

Abstract

The Marcus-Lushnikov process is a finite stochastic particle system, in which each particle is entirely characterized by its mass. Each pair of particles with masses x and y merges into a single particle at a given rate $K(x, y)$. Under certain assumptions, this process converges to the solution to Smoluchowski equation, as the number of particles increases to infinity. The Marcus-Lushnikov process gives at each time the distribution of masses of the particles present in the system, but does not retain the history of formation of the particles. In this paper, we set up a historical analogue of the Marcus-Lushnikov process (built according the rules of construction of the usual Markov-Lushnikov process) each time giving what we call the historical tree of a particle. The historical tree of a particle present in the Marcus-Lushnikov process at a given time t encodes information about the times and masses of the coagulation events that have formed that particle. We prove a law of large numbers for the empirical distribution of such historical trees. The limit is a natural measure on trees which is constructed from a solution to Smoluchowski coagulation equation.

1 Presentation of the problem

1.1 Introduction

Let $E = (0, \infty)$. Let $K : E \times E \to (0, \infty)$ be a symmetric continuous function. Let S be the set of finite integer-valued measures on $(0, \infty)$. S contains elements of the form

$$x = \sum_{i=1}^{n} m_i \delta_{y_i}$$

for $n \in \mathbb{N}$ where $y_1, \ldots, y_n > 0$ are distinct and for $i \in \{1, \ldots, n\}, m_i \in \mathbb{N}$. The Marcus-Lushnikov process with coagulation kernel K is the continuous time Markov chain on S with non-zero transition rates given by

$$q(x, x') = \begin{cases} m_i m_j K(y_i, y_j) & \text{if } i < j \\ \frac{1}{2} m_i (m_i - 1) K(y_i, y_i) & \text{if } i = j \end{cases}$$

for $x' = x + \delta_{y_i + y_j} - \delta_{y_i} - \delta_{y_j}$.

Let us give a way of constructing a Marcus-Lushnikov process $(X_t)_{t \geq 0}$. Let $[N] = \{1, \ldots, N\}$ and let $y_1, \ldots, y_N > 0$ be the masses (not necessarily distinct) associated to

---

[*] University of Cambridge, Statistical Laboratory, Centre for Mathematical Sciences Wilberforce Road, Cambridge, CB3 0WB, UK e-mail: smj45@cam.ac.uk

each particle in $[N]$. Set

$$X_0 = \sum_{i=1}^{N} \delta_{y_i}.$$

For each $i < j$ take an independent random variable T_{ij} such that T_{ij} is exponential with parameter $K(y_i, y_j)$, and define

$$T = \min_{i<j} T_{ij}.$$

Set $X_t = X_0$ for $t < T$ and

$$X_T = X_t - (\delta_{y_i} + \delta_{y_j} - \delta_{y_i + y_j})$$

if $T = T_{ij}$, then begin the construction afresh from X_T. Each pair of clusters $i < j$ coagulates at rate $K(y_i, y_j)$.

Now fix μ_0 a measure on E and take a sequence $\left(x_0^N\right)_{N \geq 0} \in S$ such that

$$\mu_0^N = N^{-1} x_0^N \to \mu_0 \tag{1}$$

weakly on E as $N \to \infty$. Let $(X_t^N)_{t \geq 0}$ be Marcus-Lushnikov with kernel $\frac{K}{N}$ starting from x_0^N. For each N, we can write

$$x_0^N = \sum_{i=1}^{m(N)} \delta_{y_i^N}$$

with $m(N) \leq N$ and $y_1^N, \ldots, y_N^{m(N)} > 0$ non necessarily distinct. Set

$$\mu_t^N = N^{-1} X_t^N.$$

Without loss of generality, for the rest of the paper, we will take $m(N) = N$. Indeed, after one step in the process above, the number of masses in the system will be less than N and so we will be exactly in the case where $m(N) < N$.

Our aim in this paper is to set up a historical analogue of the process μ_t^N and to prove that it converges to a limit measure that can be constructed from the strong solution to a generalised form of Smoluchowski's equation [1],[2](to be made precise below).

Before defining precisely this new process let us explain why it is interesting to know about the history of formation of a cluster.

The Marcus-Lushnikov process [5] describes the stochastic Markov evolution of a finite system of coalescing particles. It gives at each time the distribution in masses of the particles present in the system but does not retain any other information that the particles might contain. In other words, we lose in part the information contained in the particles that is their history. Why is it interesting to know about the history ? For instance, consider a system of N particles with associated masses $y_1, \ldots, y_N > 0$. Assume that these particles can only be of three types say either A, B or C. Allow them to coagulate according to the rules of coagulation of the Marcus-Lushnikov process. Then, the usual Marcus-Lushnikov will give us at each time the masses of the particles present in the system but will not be able to tell us for each particle present at this time how many particles of type A, B or C this particle contains along with the order of formation. Our historical measure will give us at each time the particles formed with their respective masses, the time when they have formed but also the history of the formation from its beginning that is the time at which each intermediary particles have formed along with what they contain. We could think of an other application in industry in the process of making a certain chemical product. We can assume that in order to make a certain powder we need to put N ingredients in a specific order and at specific times. Then our historical measure will allow us to follow the formation of the powder and to detect if ingredients were put in the wrong order at the wrong time.

We are now going to review the work of [1] and [2] about the convergence of μ_t^N to the strong solution to Smoluchowski's equation as we will use this tool to prove our main result (stated in 1.3).

1.2 Related work

Take $\varphi : E \to (0, \infty)$ to be a continuous sublinear function. We suppose that $\varphi \geq 1$. Assume that the coagulation kernel can be written as follows:

$$K(x, y) = \tilde{K}(x, y)\varphi(x)\varphi(y) \tag{2}$$

with $\tilde{K}$ bounded on $E \times E$. For μ a non-negative Borel measure on E such that,

$$\int_{E\times E} K(x, y)\mu(dx)\mu(dy) < \infty$$

we define $L(\mu)$ as follows:

$$\langle f, L(\mu)\rangle = \frac{1}{2}\int_{E\times E} \big(f(x + y) - f(x) - f(y)\big)K(x, y)\mu(dx)\mu(dy)$$

for all f bounded and measurable. We consider the following measure-valued form of the Smoluchowski coagulation equation,

$$\mu_t = \mu_0 + \int_0^t L(\mu_s)ds. \tag{3}$$

We admit as a strong local solution any map:

$$t \to \mu_t : [0, T) \to \mathcal{M}^+$$

where $T \in (0, \infty]$ and $\mathcal{M}^+$ is the set of non-negative Borel measures on E, such that

1. for all $B \subseteq E$ compact,

$$t \to \mu_t(B) : [0, T) \to (0, \infty)$$

is measurable.

2.

$$\int_0^t \int_E \varphi^2(x)\mu_s(dx) < \infty$$

for all $t < T$.

3. for all bounded measurable functions f, for all $t < T$,

$$\int_E f(x)\mu_t(dx) = \int_E f(x)\mu_0(dx) + \int_0^t \int_{E\times E} \big(f(x+y) - f(x) - f(y)\big)K(x,y)\mu_s(dx)\mu_s(dy)ds.$$

In the case $T = \infty$ we call a strong local solution a strong solution. Assume that

$$\langle \varphi^2, \mu_0\rangle = \int_E \varphi^2(x)\mu_0(\mathrm{d}x) < \infty. \tag{4}$$

Then [2] tells us that there exists a unique maximal strong solution to (3) denoted $(\mu_t)_{t<T}$ for some $T > 0$. Moreover assume that

$$\int_E f(x)\varphi(x)\mu_0^N(dx) \to \int_E f(x)\varphi(x)\mu_0(dx) \tag{5}$$

as $N \to \infty$ for all f bounded and continuous on E. Then, for all $t < T$, for all f continuous and bounded on E

$$\int_E f(x)\varphi(x)\mu_t^N(dx) \to \int_E f(x)\varphi(x)\mu_t(dx)$$

as $N \to \infty$ in probability, that is

$$\mu_t^N \to \mu_t$$

as $N \to \infty$ weakly in probability with weight function φ. Indeed, for $\epsilon > 0$, for all f continuous and bounded on E, for all $t < T$, we can find $\nu > 0$, such that for all $N > 0$,

$$\mathbb{P}\left(\sup_{s\le t}\left|\int_E f(x)\varphi(x)\mu_s^N(dx) - \int_E f(x)\varphi(x)\mu_s(dx)\right| > \epsilon\right) \le \exp(-\nu N). \tag{6}$$

1.3 Our main result

Assumptions

The following assumptions hold for the whole paper. Let μ_0 be a measure on $(0,\infty)$ and take a sequence $\left(\mu_0^N\right)_{N\ge 0}$ (as defined in subsection 1.1) such that (1) is satisfied that is

$$\mu_0^N = \frac{1}{N}\sum_{k=1}^N \delta_{y_k^N} \to \mu_0 \text{ weakly as } N \to \infty.$$

For convenience we will write $y_k^N = y_k$ for all $k \in [N] = \{1,\dots,N\}$. Let $E = (0,\infty)$. Let $K : E\times E \to (0,\infty)$ be a symmetric continuous function. Let $\varphi : E \to (0,\infty)$ be a continuous sublinear function. We suppose that $\varphi \ge 1$. Assume that (2),(4) and (5) are satisfied and that K is bounded on sets of the form $[0,M]^2$ with $M > 0$. Let $(\mu_t)_{t<T}$ be the strong solution to (3) with $T > 0$.

Basic notations for Trees

For a finite set $J \subset \mathbb{N}$, we write $\mathbb{T}(J)$ for the smallest set containing J, such that $\{i, j\} \in \mathbb{T}(J)$ whenever $i, j \in \mathbb{T}(J)$. We refer to elements of $\mathbb{T}(J)$ as trees. They are finite binary trees with leaves labeled by elements of J. Each $i \in \mathbb{T}(J)$ has a set of leaf labels $\lambda(i) \subseteq J$ determined by

$$\lambda(i) = \{i\}$$

for $i \in J$ and

$$\lambda(\{i, j\}) = \lambda(i) \cup \lambda(j)$$

for all $i, j \in \mathbb{T}(J)$. For each $i \in \mathbb{T}(J)$, $|\lambda(i)|$ will denote the number of elements of the set $\lambda(i)$. Let $n : \mathbb{T}(J) \to \mathbb{N}$ be the counting function, defined as follows:

$$n(i) = 1$$

for $i \in J$ and

$$n(\{i, j\}) = n(i) + n(j)$$

for all $i, j \in \mathbb{T}(J)$. Finally, for $J \subset \mathbb{N}$ finite, we define

$$\mathbb{T}^\star(J) = \mathbb{T}^\star J = \{i \in \mathbb{T}(J) : |\lambda(i)| = n(i)\}.$$

This is the set of rooted binary trees with leaves labeled by distinct elements of J.

The coagulation process on Trees

According to the rules of construction of the usual Marcus-Lushnikov process, we set up an analogue of this process, each time giving the trees present in the system. We start with the set of labeled particles $[N] = \{1, \ldots, N\}$ with masses $y_1, \ldots, y_N > 0$. For $i = \{i_1, i_2\} \in \mathbb{T}^\star[N]$, the mass y_i of i is given recursively by

$$y_i = y_{i_1} + y_{i_2}.$$

Each tree particle $i \in \mathbb{T}^\star[N]$ has a tree type $\tau(i) \in \mathbb{T}$, where $\mathbb{T} = \mathbb{T}(\{1\})$, determined by $\tau(i) = 1$ for $i \in [N]$, and $\tau(\{i, j\}) = \{\tau(i), \tau(j)\}$ for $i, j \in \mathbb{T}^\star[N]$. We construct the historical analogue of the Marcus-Lushnikov process on $\mathbb{T}^\star[N]$ as follows. Set $I[0] = [N]$. For $i \in [N]$, set $S_i = 0$. There exist trees $i[1], \ldots, i[N-1] \in \mathbb{T}^\star[N]$, coagulation times $S_{i[1]} < \cdots < S_{i[N-1]} < \infty$ (they are exactly the jump times in the usual Marcus-Lushnikov process), and subsets $I[1], \ldots, I[N-1] \subset \mathbb{T}^\star[N]$, such that for each $n \in \{1, \ldots, N-1\}$, $(\lambda(i) : i \in I[n])$ is a partition of $[N]$, and $I[n]$ satisfies the recursive relation

$$I[n] = I[n-1] \backslash \{i_1[n], i_2[n]\} \cup \{i[n]\}$$

where $i[n] = \{i_1[n], i_2[n]\}$.

Let us give an example with $N = 5$. The drawing below represents one of the configurations we can obtain from the coagulation process on trees described above.



Here, $I[1] = \Big\{\{1,2\},3,4,5\Big\}, I[2] = \Big\{\{\{1,2\},3\},4,5\Big\}, I[3] = \Big\{\{\{1,2\},3\},\{4,5\}\Big\}$ and $I[4] = \Big\{\{\{\{1,2\},3\},\{4,5\}\}\Big\}$.

The historical measures

Fix $t < T$. Our principal object of interest is a process of empirical particle measures $\tilde{\mu}_t^N$ on the space of historical trees $A(0,t)$ which we shall now define. The space $A(0,t)$ is given by

$$A(0,t) = \bigcup_{\tau \in \mathbb{T}} A_\tau(0,t)$$

where $A_1(0,t) = (0,\infty)$ and for $\tau = \{\tau_1, \tau_2\} \in \mathbb{T}$,

$$A_\tau(0,t) = \big\{(s, \{\xi_1, \xi_2\}) : s \in (0,t), \xi_1 \in A_{\tau_1}(0,s), \xi_2 \in A_{\tau_2}(0,s)\big\}.$$

Let us illustrate these historical spaces through an example. Take $\tau = \{\{1\},1\} \in \mathbb{T}$. The tree below represents a $\xi = (s_2, \{\xi_1, \xi_2\}) \in A_\tau(0,t)$ with $\xi_1 \in A_{\{1\}}(0,s_2), \xi_2 \in A_1(0,s_2)$.



Here $\xi_1 = (s_1, \{y_1, y_2\}), \xi_2 = y_3$ and $\xi = \Big(s_2, \{(s_1, \{y_1, y_2\}), y_3\}\Big)$.

We equip $A(0,t)$ with its Borel σ-algebra (we explain in Appendix 5.1 how to equip $A(0,t)$ with a topology). We define on $A(0,t)$ the mass function $m : A(0,t) \to (0,\infty)$.

For $\xi \in A_1(0,t) = (0,\infty)$, we set

$$m(\xi) = \xi.$$

Recursively for $\tau = \{\tau_1, \tau_2\} \in \mathbb{T}$, for $\xi = (s, \{\xi_1, \xi_2\}) \in A_\tau(0,t)$, we set

$$m(\xi) = m(\xi_1) + m(\xi_2).$$

The empirical historical measure $\tilde{\mu}_t^N$ is given by

$$\tilde{\mu}_t^N = \frac{1}{N} \sum_{i \in I(t)} \delta_{\xi_t^i}$$

where $I(t) \subset \mathbb{T}^\star[N]$ is the set of trees present in the system at time t. $I(t)$ is given by

$$I(t) = I[n] \text{ for } S_{i[n]} \leq t < S_{i[n+1]}.$$

For $i \in [N], \xi_t^i = y_i$ and for $i = \{i_1, i_2\} \in I(t)$, with $S_i = s$, we set $\xi_t^i = (s, \{\xi_s^{i_1}, \xi_s^{i_2}\})$. As we trace back the past of a particle we obtain a tree. Observe that this empirical measure $\tilde{\mu}_t^N$ and our usual Marcus Lushnikov process μ_t^N (defined in subsection 1.1) are related through the following equality,

$$\mu_t^N = \tilde{\mu}_t^N \circ m^{-1}.$$

We are interested in taking the limit of this empirical measure as $N \to \infty$. We define the limit measure on $A(0,t)$ as follows. For $\xi \in A_1(0,t)$, we set

$$\tilde{\mu}_t(d\xi) = \exp\left(-\int_0^t \int_E K(y,y')\mu_r(dy')dr\right)\mu_0(d\xi)$$

where $y = m(\xi)$ and $(\mu_r)_{r<T}$ is the strong deterministic solution to (3). Recursively for $\tau = \{\tau_1, \tau_2\} \in \mathbb{T}$, $\xi = (s, \{\xi_1, \xi_2\}) \in A_\tau(0,t)$ with $s < t < T$, we define

$$\tilde{\mu}_t(d\xi) = \epsilon(\tau) K(m(\xi_1), m(\xi_2)) \tilde{\mu}_s(d\xi_1)\tilde{\mu}_s(d\xi_2) \exp\left(-\int_s^t \int_E K(y,y')\mu_r(dy')\mathrm{d}r\right) ds$$

where $y = m(\xi)$, $\epsilon(\tau) = 1$ if $\tau_1 \neq \tau_2$ and $\epsilon(\tau) = \frac{1}{2}$ if $\tau_1 = \tau_2$.

Our main result

Our aim in this paper is to prove the following result.

Theorem 1.1. *For all $t < T$,*

$$\tilde{\mu}_t^N \to \tilde{\mu}_t \tag{7}$$

weakly on $A(0,t)$ in probability.

This theorem is proved is Section 4. Before giving in subsection 1.5, an outline of how we are going to prove Theorem 1.1, we need first to introduce some more material. The next subsection 1.4 is dedicated to introduce a coupled family of processes built on the same probability space $(\Omega, \mathcal{F}, \mathbb{P})$ (to be specified below), which we shall see are Marcus-Lushnikov and which will be really convenient to use for most of our intermediary proofs.

1.4 A coupled family of Marcus-Lushnikov processes

We start with the set of particles $[N]$ with associated masses $y_1, \ldots, y_N > 0$ (that is μ_0^N). For $J \subseteq [N]$ we set $\mathbb{T}_+^\star J = \mathbb{T}^\star J \backslash J$.

A probability space and some random variables

Let $\Omega = (0,\infty)^{\mathbb{T}^\star_+[N]}$. A probability measure $\mathbb{P}$ on Ω is defined for $\omega = (\omega_i : i \in \mathbb{T}^\star_+[N])$ by

$$\mathbb{P}(d\omega) = \prod_{i \in \mathbb{T}^\star_+[N]} \exp(-\omega_i) d\omega_i.$$

Let $(U_i : i \in \mathbb{T}^\star_+[N])$ be a family of random variables on Ω defined as follows. For $\omega = (\omega_i : i \in \mathbb{T}^\star_+[N]) \in \Omega$, set $U_i(\omega) = \omega_i$. Then, under $\mathbb{P}$, $(U_i : i \in \mathbb{T}^\star_+[N])$ is a family of independent exponential random variables with parameter 1. Set $\mathcal{F} = \sigma(U_i : i \in \mathbb{T}^\star_+[N])$. Let us now define on Ω a family of random variables from which we will construct our coupled family of Marcus-Lushnikov processes. For $i \in [N]$ set

$$S_i = 0.$$

For $\{i,j\} \in \mathbb{T}^\star[N]$, define recursively

$$S_{\{i,j\}} = \max(S_i, S_j) + \frac{N}{K(y_i,y_j)} U_{\{i,j\}}.$$

The coupled family of processes

We are going to build a coupled family of processes $\Big(\big(X^J_t\big)_{t\geq 0} : J \subseteq [N]\Big)$ as follows. Fix $J \subseteq [N]$ and set $n_J = |J|$. Set $\mathcal{F}_J = \sigma(U_i : i \in \mathbb{T}^\star_+ J)$ and $\Omega^J = (0,\infty)^{\mathbb{T}^\star_+ J}$. Fix $\omega = \big(\omega_i : i \in \mathbb{T}^\star_+ J\big) \in \Omega^J$. We start off with the set of particles $J^0_N = J$ with respective masses $(y_i : i \in J)$. We set

$$X^J_0 = \sum_{i \in J} \delta_{y_i}$$

and we consider

$$v^0 = \min_{\substack{i,j \in J^0_N \\ \text{with } i \neq j}} S_{\{i,j\}}(\omega).$$

Almost surely, we have $v^0 = S_{\{i_0,j_0\}}(\omega)$ for some unique $i_0, j_0 \in J^0_N$ with $i_0 \neq j_0$. We then obtain a new set of particles $J^1_N = J^0_N \backslash \{i_0, j_0\} \cup \{\{i_0, j_0\}\}$.
We set

$$X^J_t(\omega) = X^J_0$$

for $t < v^0$ and

$$X^J_{v^0}(\omega) = \sum_{i \in J^1_N} \delta_{y_i}.$$

Now, starting from J^1_N, we consider

$$v^1 = \min_{\substack{i,j \in J^1_N \\ \text{with } i \neq j}} S_{\{i,j\}}(\omega).$$

Almost surely, we have $v^1 = S_{\{i_1,j_1\}}(\omega)$ for some unique $i_1, j_1 \in J^1_N$ with $i_1 \neq j_1$. We obtain a new set of particles $J^2_N = J^1_N \backslash \{i_1, j_1\} \cup \{\{i_1, j_1\}\}$. We set

$$X^J_t(\omega) = X^J_{v^0}(\omega)$$

for $v^0 \leq t < v^1$ and

$$X_{v^1}^J(\omega) = \sum_{i \in J_N^2} \delta_{y_i}.$$

We start again as above with J_N^2 and so on. The process stops when there is only one particle left in the system.

Therefore, for each $\omega \in \Omega^J$, there exist trees $i[1], \ldots, i[n_J - 1] \in \mathbb{T}^\star J$, coagulation times $S_{i[1]} < \cdots < S_{i[n_J-1]} < \infty$ (they are some of the $S_i(\omega)'$s), and subsets $J_N^1, \ldots, J_N^{n_J-1} \subset \mathbb{T}^\star J$, such that for each $n \in \{1, \ldots, n_J - 1\}$, $(\lambda(i) : i \in J_N^n)$ is a partition of J, and J_N^n satisfies the recursive relation

$$J_N^n = J_N^{n-1} \backslash \{i_1[n], i_2[n]\} \cup \{i[n]\}$$

where $i[n] = \{i_1[n], i_2[n]\}$. We set

$$T_i^J(\omega) = \begin{cases} S_{i[n]} & \text{if } i \in J_N^{n-1} \backslash J_N^n \\ \infty & \text{if } i = i[n_J - 1] \\ S_i(\omega) & \text{if } i \notin \cup_{n=0}^{n_J-1} J_N^n \end{cases}$$

T_i^J can be interpreted as the time of death of the tree particle i in J, if this particle was alive in the system at some time. If $J = [N]$, we set $T_i^{[N]} = T_i$. The empirical historical measure on trees $\tilde{\mu}_t^N$ can be rewritten as follows

$$\tilde{\mu}_t^N = \frac{1}{N} \sum_{i \in \mathbb{T}^\star[N]} 1_{\{S_i \leq t < T_i\}} \delta_{\xi_t^i}.$$

The Marcus Lushnikov property for this family

Theorem 1.2. *Let $J \subseteq [N]$. The process $\left(X_t^J\right)_{t \geq 0}$ is Marcus-Lushnikov with kernel $\frac{K}{N}$ starting from*

$$X_0^J = \sum_{i \in J} \delta_{y_i}.$$

In particular, $\left(X_t^{[N]}\right)_{t \geq 0}$ and $\left(X_t^N\right)_{t \geq 0}$ have same distribution.

Proof of Theorem 1.2 : Starting from

$$X_0^J = \sum_{i \in J} \delta_{y_i},$$

it is clear that the first jump has the correct distribution for Marcus-Lushnikov. Let us now look at the $k + 1^{th}$ jump. We condition on

$$A_k = \left(X_0^J = \sum_{i \in J} \delta_{y_i}, X_{s_1}^J = \sum_{i \in J_N^1} \delta_{y_i}, \ldots, X_{s_k}^J = \sum_{i \in J_N^k} \delta_{y_i} \right),$$

where for $l \in \{1, \ldots, k\}$, J_N^l is the set of particles present on $[s_l, s_{l+1})$ and $s_1 < \cdots < s_k$ are the jump times. What are the transition rates to go from step k to step $k + 1$? Conditional on A_k, we consider the set of particles J_N^k and look at

$$T' = \min_{\substack{i,j \in J_N^k \\ \text{with } i \neq j}} S_{\{i,j\}} = \min_{\substack{i,j \in J_N^k \\ \text{with } i \neq j}} s_{\{i,j\}} + V_{\{i,j\}}$$

where for each $i,j \in J_N^k$ with $i \neq j$, $V_{\{i,j\}}$ is exponential with parameter $\frac{K(y_i,y_j)}{N}$ and,

$$s_{\{i,j\}} = \begin{cases} 0 & \text{if } i,j \in J\ . \\ s_i & \text{if } i \text{ was formed at some } s_i \in \{s_1,\dots,s_k\} \text{ and } j \in J. \\ s_j & \text{if } j \text{ was formed at some } s_j \in \{s_1,\dots,s_k\} \text{ and } i \in J. \\ \max(s_i,s_j) & \text{if } i \text{ was formed at some } s_i \in \{s_1,\dots,s_k\} \text{ and} \\ & j \text{ was formed at some } s_j \in \{s_1,\dots,s_k\}\backslash\{s_i\}. \end{cases}$$

Thus, for each $i,j \in J_N^k$ with $i \neq j$, $V_{\{i,j\}}$ has started at time $s_{\{i,j\}}$ and has been running for a duration of $s_k - s_{\{i,j\}}$. Nevertheless, by the memoryless property for exponential random variables, for $h > 0$,

$$\mathbb{P}\left(s_{\{i,j\}} + V_{\{i,j\}} > s_k + h | s_{\{i,j\}} + V_{\{i,j\}} > s_k\right) = \mathbb{P}\left(V_{\{i,j\}} > h\right).$$

Thus, it is equivalent to add $s_k - s_{\{i,j\}}$ to $S_{\{i,j\}}$ and consider that $V_{\{i,j\}}$ starts from s_k. Hence, conditional on A_k, we find exactly the T we considered in the construction of the the Marcus-Lushnikov process in 1.1. Therefore, the transition rates are the same. Thus $\left(X_t^J\right)_{t\geq 0}$ is Marcus-Lushnikov.

□

For all $t \geq 0$, we set $\mu_t^{N,J} = N^{-1}X_t^J$. Note that by construction, for all $J \subseteq [N]$, $(\mu_t^{N,J})_{t\geq 0}$ is measurable with respect to $\mathcal{F}_J = \sigma(U_i : i \in \mathbb{T}_+^\star J)$. We shall see that the coupled family of Marcus-Lushnikov processes $\left(\left(\mu_t^{N,J}\right)_{t\geq 0} : J \subseteq [N]\right)$ will be useful in most of the intermediary proofs leading to our main result. Let us now give an outline of the intermediary results we need in order to prove Theorem 1.1.

1.5 Outline proof of main result

Fix $t \in [0,T)$. In order to prove Theorem 1.1, we need to prove that for all $f \in C_b\left(A(0,t)\right)$

$$\langle f, \tilde{\mu}_t^N \rangle = \int_{A(0,t)} f(\xi)\tilde{\mu}_t^N(d\xi) \to \langle f, \tilde{\mu}_t \rangle = \int_{A(0,t)} f(\xi)\tilde{\mu}_t(d\xi) \tag{8}$$

as $N \to \infty$ in probability. To prove (8), we shall see that it is sufficient to prove that for all $\tau \in \mathbb{T}$, for all $f \in C_b\left(A_\tau(0,t)\right)$

$$\langle f, \tilde{\mu}_t^N \rangle = \int_{A_\tau(0,t)} f(\xi)\tilde{\mu}_t^N(d\xi) \to \langle f, \tilde{\mu}_t \rangle = \int_{A_\tau(0,t)} f(\xi)\tilde{\mu}_t(d\xi) \tag{9}$$

as $N \to \infty$ in probability. Then we will be able to conclude, using a tightness argument (that is explained in subsection 4.1). To show (9), it is sufficient to prove that

$$\mathbb{E}\left(\langle f, \tilde{\mu}_t^N \rangle\right) \to \langle f, \tilde{\mu}_t \rangle \tag{10}$$

and

$$\mathbb{E}\left(\langle f, \tilde{\mu}_t^N \rangle^2\right) \to \langle f, \tilde{\mu}_t \rangle^2 \tag{11}$$

as $N \to \infty$. Then,

$$\mathbb{E}\left(\langle f, \tilde{\mu}_t^N - \tilde{\mu}_t \rangle^2\right) \to 0$$

and a fortiori

$$\langle f, \tilde{\mu}_t^N \rangle \to \langle f, \tilde{\mu}_t \rangle$$

as $N \to \infty$ in probability.

Thus, we need to compute $\mathbb{E}\left(\langle f, \tilde{\mu}_t^N \rangle\right)$. The paper is set as follows. Section 2 is dedicated to the computation of $\mathbb{E}\left(\langle f, \tilde{\mu}_t^N \rangle\right)$. In Section 3, we will prove (10) and (11) in order to obtain (9). Finally in Section 4 we will use the results out obtained in the previous sections to prove Theorem 1.1.

2 Intermediary computations

For the whole section we start with the set of particles $[N]$ with associated masses $y_1, \dots, y_N > 0$ (that is μ_0^N) and we consider the probability space $(\Omega, \mathcal{F}, \mathbb{P})$ that we have defined in subsection 1.4. We fix $0 \le t < T$. The aim of this section is to calculate $\mathbb{E}\left(\langle f, \tilde{\mu}_t^N \rangle\right)$ for $f \in C_b(A_\tau(0,t))$ with $\tau \in \mathbb{T}$.

2.1 A finite sum of conditional expectations

Let us fix $\tau \in \mathbb{T}$ with n leaves that is $n(\tau) = n$ and take $f \in C_b(A_\tau(0,t))$. We are going to express $\mathbb{E}\left(\langle f, \tilde{\mu}_t^N \rangle\right)$ as the expectation of a finite sum of conditional expectations. Recall that for $i \in \mathbb{T}^\star[N]$, $\tau(i)$ denotes the type of the labeled tree i. We can write,

$$\begin{aligned}\langle f, \tilde{\mu}_t^N \rangle =& \frac{1}{N} \sum_{i \in \mathbb{T}^\star[N]} f\left(\xi_t^i\right) 1_{\{S_i \le t < T_i\}} \\ =& \frac{1}{N} \sum_{\substack{i \in \mathbb{T}^\star[N] \\ \text{with } \tau(i) = \tau}} f\left(\xi_t^i\right) 1_{\{S_i \le t < T_i\}}\end{aligned}$$

as f is supported on $A_\tau(0,t)$. For $J \subseteq [N]$, and $k \in \mathbb{N}$ define

$$\mathbb{T}_k^\star J = \{i \in \mathbb{T}^\star J : n(i) = k\}.$$

For any $i \in \mathbb{T}_n^\star[N]$ with type τ, there are $2^{q(\tau)}$ permutations possible of the particles composing i which will keep the tree i invariant where $q(\tau)$ is the number of symmetries in the tree τ. This is given recursively by $q(1) = 0$ and for $\tau = \{\tau_1, \tau_2\} \in \mathbb{T}$, $q(\tau) = q(\tau_1) + q(\tau_2) + 1_{\{\tau_1 = \tau_2\}}$.

Example : Take $i = \{1, \{2, 3\}\}$. This tree has for type $\tau = \{1, \{1\}\}$ and $q(\tau) = 1$. The permutations leaving the tree i alike are the identity permutation and the one sending 1 to itself and exchanging 2 and 3. Hence $2^{q(\tau)}$ permutations leave this tree invariant.

Now fix $i_0 \in \mathbb{T}_n^\star[n]$ with type $\tau(i_0) = \tau$. Define

$$S_{n,N} = \Big\{ \text{ injections } \sigma : [n] \to [N] \Big\}.$$

For $\sigma \in S_{n,N}$, set $\sigma(1,\dots,n) = \big(\sigma(1),\dots,\sigma(n)\big)$. Let

$$[N]^{\star}_{[n]} = \Big\{(i_1,\dots,i_n) \in [N]^n : i_1,\dots,i_n \text{ distinct}\Big\}.$$

Observe that for each $(i_1,\dots,i_n) \in [N]^{\star}_{[n]}$ there is a unique $\sigma \in S_{n,N}$ such that $\sigma(1,\dots,n) = (i_1,\dots,i_n)$. Hence,

$$[N]^{\star}_{[n]} = \Big\{\sigma(1,\dots,n) : \sigma \in S_{n,N}\Big\}.$$

For $\sigma \in S_{n,N}$ define $\sigma(i_0) \in \mathbb{T}^{\star}_n[N]$ to be the tree obtained from i_0 by replacing each particle $i \in [n]$ in i_0 by $\sigma(i)$.

Example : If $N = 5$, $i_0 = \{1,\{2,3\}\}$ and $\sigma \in S_{3,5}$ is such that $\sigma(1) = 3, \sigma(2) = 5, \sigma(3) = 1$, then

$$\sigma(i_0) = \{3,\{5,1\}\}.$$

Define

$$I_0 : [N]^{\star}_{[n]} \to \mathbb{T}^{\star}_n[N] \text{ by } I_0\left(\sigma(1,\dots,n)\right) = \sigma(i_0).$$

Thus, we can write

$$\begin{aligned}
\langle f, \tilde{\mu}^N_t \rangle &= \frac{2^{-q(\tau)}}{N} \sum_{\sigma \in S_{n,N}} f\left(\xi^{\sigma(i_0)}_t\right) 1_{\{S_{\sigma(i_0)} \le t < T_{\sigma(i_0)}\}} \\
&= \frac{2^{-q(\tau)}}{N} \sum_{(i_1,\dots,i_n) \in [N]^{\star}_{[n]}} f\left(\xi^{I_0(i_1,\dots,i_n)}_t\right) 1_{\{S_{I_0(i_1,\dots,i_n)} \le t < T_{I_0(i_1,\dots,i_n)}\}}.
\end{aligned}$$

Hence, setting $J(\sigma(i_0)) = [N] \backslash \{\lambda(\sigma(i_0))\}$, we can write

$$\begin{aligned}
\mathbb{E}\left(\langle f, \tilde{\mu}^N_t \rangle\right) &= \frac{2^{-q(\tau)}}{N} \sum_{\sigma \in S_{n,N}} \mathbb{E}\Bigg(f\left(\xi^{\sigma(i_0)}_t\right) 1_{\{S_{\sigma(i_0)} \le t < T_{\sigma(i_0)}\}} \Bigg) \\
&= \mathbb{E}\left(\frac{2^{-q(\tau)}}{N} \sum_{\sigma \in S_{n,N}} \mathbb{E}\Bigg(f\left(\xi^{\sigma(i_0)}_t\right) 1_{\{S_{\sigma(i_0)} \le t < T_{\sigma(i_0)}\}} | \mathcal{F}^{J(\sigma(i_0))} \Bigg) \right).
\end{aligned}$$

We shall compute

$$\mathbb{E}\Bigg(f\left(\xi^{\sigma(i_0)}_t\right) 1_{\{S_{\sigma(i_0)} \le t < T_{\sigma(i_0)}\}} | \mathcal{F}^{J(\sigma(i_0))} \Bigg) \tag{12}$$

for all $\sigma \in S_{n,N}$. In order to compute this quantity, we shall find it useful to work with labeled trees.

2.2 Working with labeled trees

We are going to introduce spaces similar to $A(0,t)$, but for labeled trees.

Vector tree of masses

Each particle $j \in [N]$ has a given mass y_j. Take a tree $i \in \mathbb{T}^\star_n[N]$. Assume that this tree is formed from the particles $i_1 < \cdots < i_n$. We write $y = (y_{i_1}, \ldots, y_{i_n})$ for its associated vector of masses. Its vector tree of masses $\tilde{y}$ is the tree of masses obtained from i by replacing each particle in this tree by its mass. For instance $i = \{1, \{2, 3\}\}$ and $y = (y_1, y_2, y_3)$ give $\tilde{y} = \{y_1, \{y_2, y_3\}\}$.

Labeled historical spaces of trees

Define $A^{y_k}_k(0,t)$ for $k \in \{1, \ldots, N\}$, by

$$A^{y_k}_k(0,t) = \{k\} \times \{y_k\}$$

and recursively for $i = \{i_1, i_2\} \in \mathbb{T}^\star_n[N]$ define

$$A^y_i(0,t) = \left\{\, (s, \{\xi_1, \xi_2\}) : s \in (0,t), \xi_1 \in A^{y^1}_{i_1}(0,s), \xi_2 \in A^{y^2}_{i_2}(0,s) \right\}$$

where y, y^1 and y^2 are the respective vectors of masses of i, i_1 and i_2. For $i \in \mathbb{T}^\star_n[N]$ (without any associated masses) define

$$A_i(0,t) = \bigcup_{y \in (0,\infty)^n} A^y_i(0,t)$$

and set

$$\tilde{A}(0,t) = \bigcup_{N=1}^{\infty} \bigcup_{i \in \mathbb{T}^\star[N]} A_i(0,t).$$

Observe that when we integrate over $A^y_i(0,t)$, we only integrate over the coagulation times as the masses of the particles are fixed whereas integrating over $A_i(0,t)$ means integrating over the coagulation times along with the masses.

Reduction of the problem

Each particle $j \in [N]$ has a given mass y_j. Take $\tau \in \mathbb{T}$ with $n(\tau) = n$ for some $n \in \mathbb{N}$. Fix $f \in C_b(A_\tau(0,t))$. In order to compute $\mathbb{E}\left(\langle f, \tilde{\mu}^N_t \rangle\right)$, without loss of generality, by subsection 2.1, we need to compute

$$\mathbb{E}\left(f\left(\xi^i_t\right) 1_{\{S_i \le t < T_i\}} | \mathcal{F}^J \right) N^{n-1}$$

for $i \in \mathbb{T}^\star_n[n]$ with type $\tau(i) = \tau$ and associated vector of masses $y = (y_1, \ldots, y_n)$ and $J = [N] \backslash [n]$. The map

$$g_i : A_i(0,t) \to A_\tau(0,t)$$

on forgetting labels is $2^{q(\tau)}$ to 1. Define

$$f_i = f \circ g_i.$$

We have $f_i \in C_b\left(A_i(0,t)\right)$. Let $\xi_t^i \in A_\tau(0,t)$. Then,

$$g_i^{-1}(\xi_t^i) = \Big\{\zeta_t \in A_i(0,t) : g_i(\zeta_t) = \xi_t^i\Big\}.$$

The set $g_i^{-1}(\xi_t^i)$ contains $2^{q(\tau)}$ elements. Hence, we can write

$$f\left(\xi_t^i\right) = 2^{-q(\tau)} \sum_{\zeta_t \in g_i^{-1}(\xi_t^i)} f\left(g_i(\zeta_t)\right).$$

So,

$$\mathbb{E}\Big(f\left(\xi_t^i\right) 1_{\{S_i \le t < T_i\}}\Big) = 2^{-q(\tau)} \mathbb{E}\Big(\sum_{\zeta_t \in g_i^{-1}(\xi_t^i)} f\left(g_i(\zeta_t)\right) 1_{\{S_i \le t < T_i\}}\Big).$$

If we fix $\zeta_t^0 \in g_i^{-1}(\xi_t^i)$, then all the other elements of $g_i^{-1}(\xi_t^i)$ can be obtained from ζ_t^0 by permuting masses between the particles that form a symmetry in the tree i. Hence $g_i^{-1}(\xi_t^i)$ can be written as a set depending only on ζ_t^0.

Example : Take $i = \{1,2\}$. Then $\tau(i) = \tau = \{1\}$. Let $\xi_t^i = (s, \{y, y'\}) \in A_\tau(0,t)$. Then,

$$g_i^{-1}(\xi_t^i) = \Big\{(s, \{\{1\} \times \{y\}, \{2\} \times \{y'\}\}), (s, \{\{1\} \times \{y'\}, \{2\} \times \{y\}\})\Big\}.$$

It is clear in the set $g_i^{-1}(\xi_t^i)$ that we can obtain one particle from the other by exchanging masses between particles 1 and 2 which are symmetric in this tree.

Therefore,

$$\mathbb{E}\Big(f\left(\xi_t^i\right) 1_{\{S_i \le t < T_i\}}\Big) = \mathbb{E}\Big(f_i\left(\zeta_t^0\right) 1_{\{S_i \le t < T_i\}}\Big).$$

Hence,

$$\mathbb{E}\bigg(\mathbb{E}\Big(f\left(\xi_t^i\right) 1_{\{S_i \le t < T_i\}} | \mathcal{F}^J\Big)\bigg) = \mathbb{E}\bigg(\mathbb{E}\Big(f_i\left(\zeta_t^0\right) 1_{\{S_i \le t < T_i\}} | \mathcal{F}^J\Big)\bigg). \tag{13}$$

For convenience, according to the context and the spaces we consider, ξ_t^i will stand for either an element of $A_\tau(0,t)$ or $A_i(0,t)$ and more particularly since the masses of the particles are fixed as an element of $A_i^y(0,t)$ where y is the vector of masses of the tree i.

Hence it is enough to compute

$$\mathbb{E}\left(f_i\left(\xi_t^i\right) 1_{\{S_i \le t < T_i\}} | \mathcal{F}_J\right). \tag{14}$$

We are going to compute (14) first for some particular i and f_i. Then, we will use these intermediary results to solve the general case $i \in \mathbb{T}_n^\star[n]$ and $f_i \in C_b\left(A_i(0,t)\right)$ and by the relation (13) we will obtain an expression for $f \in C_b\left(A_\tau(0,t)\right)$.

2.3 Case $f = 1$ and $\tau = 1$

We take $\tau = 1 \in \mathbb{T}$ and $f = 1 \in C_b\left(A_\tau(0,t)\right)$. The corresponding set of labeled trees with type 1 is $[N]$. Without loss of generality take $i = 1$. The corresponding $f_1 \in C_b\left(A_i(0,t)\right)$ defined in subsection 2.2 is $f_1 = 1$. Also, since $i \in [N]$, we have by definition $S_i = 0$. Hence, in this case, we want to compute $\mathbb{P}(T_1 > t \mid \mathcal{F}_{J_1})$ where $J_1 = \{2, \ldots, N\}$.

Theorem 2.1. *Let* $J_1 = \{2, \ldots, N\}$ *and* $\mathcal{F}_{J_1} = \sigma\left(U_i : i \in \mathbb{T}^\star_+ J_1\right)$. *Then,*

$$\mathbb{P}(T_1 > t \mid \mathcal{F}_{J_1}) = \exp\left(-\int_0^t \int_E K(y_1, y)\mu_r^{N,J_1}(dy)dr\right) \ a.s$$

where T_1 *is as defined previously the time at which particle* 1 *dies in* $(\mu_r^N)_{r<t}$ *and* y_1 *is the mass of particle* 1*.*

Proof of Theorem 2.1 : Since μ_t^{N,J_1} is measurable with respect to $\mathcal{F}_{J_1}$, we can write $\mu_t^{N,J_1} = F_t^N(U_i : i \in \mathbb{T}^\star_+ J_1)$ where $F_t^N : (0,\infty)^{\mathbb{T}^\star_+ J_1} \to (0,\infty)$ is a measurable function. Fix $\omega^{J_1} = (\omega_i^{J_1} : i \in \mathbb{T}^\star_+ J_1) \in (0,\infty)^{\mathbb{T}^\star_+ J_1}$ and set $\mu_t^{N,J_1}(\omega^{J_1}) = F_t^N(\omega_i^{J_1} : i \in \mathbb{T}^\star_+ J_1)$ (this notation will be kept for the whole paper).

There exist $r_1(\omega^{J_1}) < \cdots < r_n(\omega^{J_1})$ such that $\mu_r^{N,J_1}(\omega^{J_1})$ is constant on the interval $[r_k(\omega^{J_1}), r_{k+1}(\omega^{J_1}))$ for $k = 0, \ldots, n$ with the convention $r_0(\omega^{J_1}) = 0$ and $r_{n+1}(\omega^{J_1}) = t$.

Moreover, we can write

$$T_1 = G_t^N\Big((U_{\{1,i\}} : i \in \mathbb{T}^\star J_1), (U_i : i \in \mathbb{T}^\star_+ J_1)\Big)$$

where $G_t^N : (0,\infty)^{\mathbb{T}^\star J_1} \times (0,\infty)^{\mathbb{T}^\star_+ J_1} \to \mathbb{R}$ is a measurable function. Hence,

$$T_1(\omega^{J_1}) = G_t^N\Big((U_{\{1,i\}} : i \in \mathbb{T}^\star J_1), \omega^{J_1}\Big)$$

is a random variable. To obtain $\mathbb{P}(T_1 > t \mid \mathcal{F}_{J_1})$, by Fubini, it is enough to compute $\mathbb{P}(T_1(\omega^{J_1}) > t)$.

Now observe that

$$\{T_1(\omega^{J_1}) > t\} = \{T_1(\omega^{J_1}) > t\} \cap \{T_1(\omega^{J_1}) > r_n(\omega^{J_1})\} \cap \ldots \cap \{T_1(\omega^{J_1}) > r_1(\omega^{J_1})\}$$

because

$$\{T_1(\omega^{J_1}) > t\} \subseteq \{T_1(\omega^{J_1}) > r_n(\omega^{J_1})\} \subseteq \ldots \subseteq \{T_1(\omega^{J_1}) > r_1(\omega^{J_1})\}.$$

Hence,

$$\mathbb{P}\left(T_1(\omega^{J_1}) > t\right) = \prod_{k=0}^{n} \mathbb{P}\left(T_1(\omega^{J_1}) > r_{k+1}(\omega^{J_1}) | T_1(\omega^{J_1}) > r_k(\omega^{J_1})\right). \tag{15}$$

Let us start by computing the quantity $\mathbb{P}\left(T_1(\omega^{J_1}) > r_1(\omega^{J_1})\right)$. Then, we will calculate $\mathbb{P}\left(T_1(\omega^{J_1}) > r_{k+1}(\omega^{J_1}) | T_1(\omega^{J_1}) > r_k(\omega^{J_1})\right)$ for $k \in \{1, \ldots, n\}$.

For $\{i,j\} \in \mathbb{T}^\star[N]$ set

$$V_{\{i,j\}} = \frac{N}{K(y_i, y_j)} U_{\{i,j\}}.$$

Observe that $\{T_1(\omega^{J_1}) > r_1(\omega^{J_1})\} = \{S_{\{1,j\}}(\omega^{J_1}) > r_1(\omega^{J_1}) : j = 2, \dots, N\}$ and that for $j \in \{2, \dots, N\}$, $S_{\{1,j\}}(\omega^{J_1}) = V_{\{1,j\}}$. Hence,

$$\begin{aligned}\mathbb{P}\left(T_1(\omega^{J_1}) > r_1(\omega^{J_1})\right) =& \mathbb{P}\left(V_{\{1,j\}} > r_1(\omega^{J_1}) : j = 2, \dots, N\right) \\ =& \prod_{j=2}^{N} \exp\left(-\frac{K(y_1, y)}{N} r_1(\omega^{J_1})\right) \\ =& \exp\left(-\sum_{j=2}^{N} \frac{K(y_1, y)}{N} r_1(\omega^{J_1})\right)\end{aligned}$$

by independence of the $(V_{\{1,j\}} : j \in J_1)$. Now, for all $r \in [0, r_1(\omega^{J_1}))$,

$$\mu_r^{N,J_1}(\omega^{J_1}) = \frac{1}{N} \sum_{j=2}^{N} \delta_{y_j}.$$

So,

$$\int_0^{r_1(\omega^{J_1})} \int_E K(y_1, y) \mu_r^{N,J_1}(\omega^{J_1})(dy)dr = \sum_{j=2}^{N} \frac{K(y_1, y_j)}{N} r_1(\omega^{J_1}).$$

Hence ,

$$\mathbb{P}\left(T_1(\omega^{J_1}) > r_1(\omega^{J_1})\right) = \exp\left(-\int_0^{r_1(\omega^{J_1})} \int_E K(y_1, y) \mu_r^{N,J_1}(\omega^{J_1})(dy)dr\right).$$

Now, let us compute $\mathbb{P}\left(T_1(\omega^{J_1}) > r_{k+1}(\omega^{J_1}) \mid T_1(\omega^{J_1}) > r_k(\omega^{J_1})\right)$ for $k \in \{1, \dots, n\}$. For $k \in \{1, \dots, n\}$, let $J_k(\omega^{J_1})$ be the set of particles present in $\left(\mu_r^{N,J_1}(\omega^{J_1})\right)$ for $r \in [r_k(\omega^{J_1}), r_{k+1}(\omega^{J_1}))$ and assume that j_k is the particle formed at time $r_k(\omega^{J_1})$. We can write $j_k = \{j_k(1), j_k(2)\}$ with $j_k(1), j_k(2) \in J_{k-1}(\omega^{J_1})$. Observe that $S_{j_k}(\omega^{J_1}) = r_k(\omega^{J_1})$. Then,

$$\begin{aligned}&\mathbb{P}\left(T_1(\omega^{J_1}) > r_{k+1}(\omega^{J_1}) | T_1(\omega^{J_1}) > r_k(\omega^{J_1})\right) \\ &\quad = \mathbb{P}\left(S_{\{1,j\}}(\omega^{J_1}) > r_{k+1}(\omega^{J_1}) \text{ for all } j \in J_k(\omega^{J_1}) | T_1(\omega^{J_1}) > r_k(\omega^{J_1})\right).\end{aligned}$$

Observe that $J_k(\omega^{J_1}) = \{j_k\} \cup J_{k-1}(\omega^{J_1}) \backslash \{j_k(1), j_k(2)\}$. By the memoryless property for exponential random variables,

$$\begin{aligned}&\mathbb{P}\left(T_1(\omega^{J_1}) > r_{k+1}(\omega^{J_1}) \mid T_1(\omega^{J_1}) > r_k(\omega^{J_1})\right) \\ &\ = \mathbb{P}\big(S_{\{1,j\}}(\omega^{J_1}) > r_{k+1}(\omega^{J_1}) \text{ for all } j \in J_k(\omega^{J_1}) \backslash \{j_k\}, \\ &\qquad S_{\{1,j_k\}}(\omega^{J_1}) > r_{k+1}(\omega^{J_1}) \mid S_{\{1,j\}}(\omega^{J_1}) > r_k(\omega^{J_1}) \text{ for all } j \in J_k(\omega^{J_1}) \backslash \{j_k\}\big).\end{aligned}$$

Now we can write $J_k(\omega^{J_1}) \backslash \{j_k\} = I_k(\omega^{J_1}) \cup K_k(\omega^{J_1})$ where $K_k(\omega^{J_1}) \subset \{j_1, \dots, j_{k-1}\}$ and $I_k(\omega^{J_1}) \subset \{2, \dots, N\}$. For $j \in I_k(\omega^{J_1})$, we have $S_{\{1,j\}}(\omega^{J_1}) = V_{\{1,j\}}$, and for $j \in K_k(\omega^{J_1})$, there exist $r_{k_j} \in \{r_1(\omega^{J_1}), \dots, r_{k-1}(\omega^{J_1})\}$ such that $S_j(\omega^{J_1}) = r_{k_j}$. So, $S_{\{1,j\}}(\omega^{J_1}) = S_j(\omega^{J_1}) + V_{\{1,j\}} = r_{k_j} + V_{\{1,j\}}$. Hence, for each $j \in J_k(\omega^{J_1}) \backslash \{j_k\}$, $S_{\{1,j\}}$ only depends on $V_{\{1,j\}}$. Also, $S_{j_k}(\omega^{J_1}) = r_k(\omega^{J_1})$ so $S_{\{1,j_k\}}(\omega^{J_1}) = r_k(\omega^{J_1}) + V_{\{1,j_k\}}$. Thus,

using the independence of $\left(V_{\{1,j\}}, j \in J_k(\omega^{J_1})\backslash\{j_k\}\right)$ and the memoryless property for exponential random variables, we obtain that :

$$\begin{aligned}
&\mathbb{P}(T_1(\omega^{J_1}) > r_{k+1}(\omega^{J_1}) \mid T_1(\omega^{J_1}) > r_k(\omega^{J_1})) \\
&\quad = \mathbb{P}\left(V_{\{1,j_k\}} > r_{k+1}(\omega^{J_1}) - r_k(\omega^{J_1})\right) \\
&\quad \prod_{j \in I_k(\omega^{J_1})} \mathbb{P}\left(V_{\{1,j\}} > r_{k+1}(\omega^{J_1}) \mid V_{\{1,j\}} > r_k(\omega^{J_1})\right) \\
&\quad \prod_{j \in K_k(\omega^{J_1})} \mathbb{P}\left(V_{\{1,j\}} > r_{k+1}(\omega^{J_1}) - r_{k_j} \mid V_{\{1,j\}} > r_k(\omega^{J_1}) - r_{k_j}\right) \\
&\quad = \mathbb{P}\left(V_{\{1,j_k\}} > r_{k+1}(\omega^{J_1}) - r_k(\omega^{J_1})\right) \\
&\quad \prod_{j \in I_k(\omega^{J_1})} \mathbb{P}\left(V_{\{1,j\}} > r_{k+1}(\omega^{J_1}) - r_k(\omega^{J_1})\right) \\
&\quad \prod_{j \in K_k(\omega^{J_1})} \mathbb{P}\left(V_{\{1,j\}} > r_{k+1}(\omega^{J_1}) - r_k(\omega^{J_1})\right).
\end{aligned}$$

Hence,

$$\begin{aligned}
&\mathbb{P}(T_1(\omega^{J_1}) > r_{k+1}(\omega^{J_1}) \mid T_1(\omega^{J_1}) > r_k(\omega^{J_1})) \\
&\quad = \prod_{j \in J_k(\omega^{J_1})} \mathbb{P}\left(V_{\{1,j\}} > r_{k+1}(\omega^{J_1}) - r_k(\omega^{J_1})\right) \\
&\quad = \prod_{j \in J_k(\omega^{J_1})} \exp\left(-\frac{K(y_1, y_j)}{N}(r_{k+1}(\omega^{J_1}) - r_k(\omega^{J_1}))\right).
\end{aligned}$$

Now, for all $r \in [r_k(\omega^{J_1}), r_{k+1}(\omega^{J_1}))$,

$$\mu_r^{N,J_1}(\omega^{J_1}) = \frac{1}{N} \sum_{j \in J_k(\omega^{J_1})} \delta_{y_j}.$$

So,

$$\int_{r_k(\omega^{J_1})}^{r_{k+1}(\omega^{J_1})} \int_E K(y_1, y) \mu_r^{N,J_1}(\omega^{J_1})(dy) dr = \sum_{j \in J_k(\omega^{J_1})} \frac{K(y_1, y_j)}{N} \left(r_{k+1}(\omega^{J_1}) - r_k(\omega^{J_1})\right).$$

Hence,

$$\begin{aligned}
&\mathbb{P}(T_1(\omega^{J_1}) > r_{k+1}(\omega^{J_1}) \mid T_1(\omega^{J_1}) > r_k(\omega^{J_1})) \\
&\quad = \exp\left(-\int_{r_k(\omega^{J_1})}^{r_{k+1}(\omega^{J_1})} \int_E K(y_1, y) \mu_r^{N,J_1}(\omega^{J_1})(dy) dr\right).
\end{aligned}$$

and using the equality (15) we obtain :

$$\mathbb{P}(T_1(\omega^{J_1}) > t) = \exp\left(-\int_0^t \int_E K(y_1, y) \mu_r^{N,J_1}(\omega^{J_1})(dy) dr\right).$$

Finally, by Fubini, we get :

$$\mathbb{P}(T_1 > t \mid \mathcal{F}_{J_1}) = \exp\left(-\int_0^t \int_E K(y_1, y)\mu_r^{N,J_1}(\mathrm{d}y)\mathrm{d}r\right).$$

as required.

$\square$

2.4 Case f and τ general

Recall that each particle $j \in [N]$ has a given mass $y_j > 0$. Take $\tau \in \mathbb{T}$ with $n(\tau) = n$, for some $n \in \mathbb{N}$ and fix $i \in \mathbb{T}_n^\star[n]$ with type $\tau(i) = \tau$ and associated vector of masses $y = (y_1, \ldots, y_n)$. Fix $f \in C_b\left(A_\tau(0,t)\right)$ and consider $f_i \in C_b(A_i(0,t))$ constructed from f as in subsection 2.2. Let $J = [N]\backslash[n]$ and set $\mathcal{F}_J = \sigma(U_j : j \in \mathbb{T}_+^\star J)$. Our aim in this subsection is to compute

$$\mathbb{E}_{y,\mu_0^{N,J}}\left(f_i\left(\xi_t^i\right)1_{\{S_i \leq t < T_i\}} | \mathcal{F}_J\right)$$

where T_i, the time of death of the labeled tree i, has been defined in the subsection 1.4. The notation "$\mathbb{E}_{y,\mu_0^{N,J}}$" means that the expectation is taken conditional on starting with the set of particles $[N]$ with respective masses $y_1, \ldots, y_N > 0$. This condition can be rewritten as follows : $y = (y_1, \ldots, y_n)$ and $\mu_0^{N,J} = \frac{1}{N}\sum_{j \in J}\delta_{y_j}$.

2.4.1 Useful notations

Fix $0 \leq t < T$.

The space $\Delta(\xi)$

For $k \in [N]$, for $\xi \in A_k^{y_k}(0,t)$, define

$$\Delta(\xi) = (0,t) \times \{\xi\}.$$

Recursively for $i \in \mathbb{T}_n^\star[n]$ with associated vector of masses y, for $\xi = (s, \{\xi_1, \xi_2\}) \in A_i^y(0,t)$, define

$$\Delta(\xi) = \Delta(\xi_1) \cup \Delta(\xi_2) \cup \big([s,t) \cup \{\xi\}\big).$$

The projection map $\Delta(\xi) \to (0,t)$

For $k \in [N]$, for $\xi \in A_k^{y_k}(0,t)$, define the projection map $\Pi = \Pi_\xi : \Delta(\xi) \to (0,t)$ by for $s \in (0,t)$,

$$\Pi_\xi\left((s, \{\xi\})\right) = s.$$

Recursively for $i \in \mathbb{T}_n^\star[n]$ with associated vector of masses y, for $\xi = (s, \{\xi_1, \xi_2\}) \in A_i^y(0,t)$, define the projection map $\Pi = \Pi_\xi : \Delta(\xi) \to (0,t)$ by

$$\Pi(u) = \Pi_\xi(u) = \begin{cases} \Pi_{\xi_1}(u) & \text{if } u \in \Delta(\xi_1) \\ \Pi_{\xi_2}(u) & \text{if } u \in \Delta(\xi_2) \\ r & \text{if } u = (r,\xi) \in [s,t) \cup \{\xi\} \end{cases}$$

The projection map from $\Delta(\xi)$ onto the space of trees of masses

For $k \in [N]$, for $\xi \in A_k^{y_k}(0,t)$, set $y_t(\xi) = y_k$ and define

$$y = y(\xi) : \Delta(\xi) \to (0,\infty) \text{ by } y(\xi)(u) = y_u = y_k.$$

Recursively for $i \in \mathbb{T}_n^\star[n]$ with associated vector of masses y, for $\xi = (s, \{\xi_1, \xi_2\}) \in A_i^y(0,t)$, set

$$y_t(\xi) = \{y_s(\xi_1), y_s(\xi_2)\}$$

and define

$$y = y(\xi) : \Delta(\xi) \to \mathbb{T}(0,\infty)$$

where the space $\mathbb{T}(0,\infty)$ is given by

$$\mathbb{T}(0,\infty) = \bigcup_{\tau \in \mathbb{T}} \mathbb{T}_\tau(0,\infty)$$

where $\mathbb{T}_1(0,\infty) = (0,\infty)$ and for $\tau = \{\tau_1, \tau_2\} \in \mathbb{T}$,

$$\mathbb{T}_\tau(0,\infty) = \big\{ y = \{y^1, y^2\} : y_1 \in \mathbb{T}_{\tau_1}(0,\infty), y_2 \in \mathbb{T}_{\tau_2}(0,\infty) \big\},$$

by

$$y(\xi)(u) = \begin{cases} y(\xi_1)(u) & \text{if } u \in \Delta(\xi_1) \\ y(\xi_2)(u) & \text{if } u \in \Delta(\xi_2) \\ y_u = \{y_s(\xi_1), y_s(\xi_2)\} & \text{if } u = (r,\xi) \in [s,t) \cup \{\xi\} \end{cases}$$

The notation K_ξ

For $\xi \in A_k(0,t)$ with $k \in [N]$, set

$$K_\xi = 1.$$

Recursively for $i \in \mathbb{T}^\star[N]$, for $\xi = (s, \{\xi_1, \xi_2\}) \in A_i(0,t)$, define

$$K_\xi = K(m(\xi_1), m(\xi_2)) K_{\xi_1} K_{\xi_2}.$$

2.4.2 An expression for the conditional expectation

Set $\mathcal{F}_{[n]} = \sigma(U_j : j \in \mathbb{T}_+^\star[n])$. The aim is to compute

$$\begin{aligned} &\mathbb{E}_{y,\mu_0^{N,J}} \left(f_i\left(\xi_t^i\right) 1_{\{S_i \le t < T_i\}} | \mathcal{F}_J \vee \mathcal{F}_{[n]} \right) \\ &\qquad = f_i\left(\xi_t^i\right) 1_{\{S_i \le t < T_i^{[n]}\}} \mathbb{E}_{y,\mu_0^{N,J}} \left(1_{\{t < T_i\}} | \mathcal{F}_J \vee \mathcal{F}_{[n]} \right) \end{aligned}$$

because $f_i\left(\xi_t^i\right) 1_{\{S_i \le t < T_i^{[n]}\}}$ is measurable with respect to $\mathcal{F}_{[n]}$ and $\{t < T_i\} \subset \{t < T_i^{[n]}\}$ ($T_i^{[n]}$ is defined in 1.4). The lemma below will be proved later on.

Lemma 2.2. *With the notations above, on the event $\{S_i \le t < T_i^{[n]}\}$,*

$$\mathbb{P}_{y,\mu_0^{N,J}} \left(t < T_i | \mathcal{F}_J \vee \mathcal{F}_{[n]} \right) = \exp\left(- \int_{\Delta(\xi_t^i)} \int_E K(y_r, y') \mu^{N,J}_{\Pi_{\xi_t^i}(r)}(dy') dr \right) \ a.s.$$

Hence, integrating over $A_i^y(0,t)$ we obtain

$$\mathbb{E}_{y,\mu_0^{N,J}}\left(f_i\left(\xi_t^i\right)1_{\{S_i\le t<T_i\}}|\mathcal{F}_J\right)=\int_{A_i^y(0,t)} f_i\left(\xi\right)\mathbb{P}_y\left(S_i\le t<T_i^{[n]},\xi_t^i\in d\xi\right)$$
$$\exp\left(-\int_{\Delta(\xi)}\int_E K(y_r,y')\mu^{N,J}_{\Pi_\xi(r)}(dy')dr\right).$$

Therefore, we need to compute $\mathbb{P}_y\left(S_i\le t<T_i^{[n]},\xi_t^i\in d\xi\right)$ for some $\xi\in A_i^y(0,t)$.

Lemma 2.3.

$$\mathbb{P}_y\left(S_i\le t<T_i^{[n]},\xi_t^i\in d\xi\right)=\frac{K_\xi}{N^{n-1}}\exp\left(-\int_0^t\frac{K_s(\xi)}{N}ds\right)\nu(d\xi)$$

where $\nu=\nu_i^y$ *is the Lebesgue measure defined in the Appendix* 5.2 *and for* $\xi\in A_i^y(0,t)$ *and* $s\in(0,t)$,

$$K_s(\xi)=\frac{1}{2}\sum_{\substack{r,r'\in\Pi_\xi^{-1}(s)\\ r\neq r'}}K(y_r,y_{r'})$$

with $\Pi_\xi:\Delta(\xi)\to(0,t)$ *as defined in 2.4.1.*

Proof of Lemma 2.3: Let us fix $\xi\in A_i^y(0,t)$. To ξ we can uniquely associate $\{(s_1,j_1),\dots,(s_{n-1},j_{n-1})\}$ where $s_1<\dots<s_{n-1}$ are the coagulation times and $j_1,\dots,j_{n-1}$ are the labeled trees (subtrees of i) formed at $s_1,\dots,s_{n-1}$ respectively. Denote by $J^k(\xi)$ the set of trees (subtrees of i) present on $[s_k,s_{k+1})$ for $k\in\{0,\dots,n-1\}$ with the convention $s_0=0$ and $s_n=t$. For each $k\in\{1,\dots,n-1\}$, we write $j_k=\{j_k(1),j_k(2)\}$ with $j_k(1),j_k(2)\in J^{k-1}(\xi)$. Observe that $J^k(\xi)=\{j_k\}\cup J^{k-1}(\xi)\backslash\{j_k(1),j_k(2)\}$. For convenience, for each $k\in\{1,\dots,n-1\}$ we will write

$$K_{j_k}=K(y_{j_k(1)},y_{j_k(2)}).$$

We start with

$$\mu_0^{N,[n]}=\frac{1}{N}\sum_{j=1}^n\delta_{y_j}.$$

Let $(Y_k)_{k=1,\dots,n-1}$ be the jump chain associated to the Markov-Lushnikov process $\left(\mu_r^{N,[n]}\right)_{r<t}$. Let $J_1,\dots,J_{n-1}$ be the jump times. Observe that after the $n-1^{th}$ jump there is only one particle left in the system, and so the process is in an absorbing state. Thus, the configuration ξ can be represented through the jump times and the jump states by, for $k=0,\dots,n-1$,

$$Y_k=\frac{1}{N}\sum_{j\in J^k(\xi)}\delta_{y_j}\text{ and }J_k=s_k.$$

For $k\in\{0,\dots,n-2\}$, the rate of going from $Y_k=\frac{1}{N}\sum_{j\in J^k(\xi)}\delta_{y_j}$ to $Y_{k+1}=\frac{1}{N}\sum_{j\in J^{k+1}(\xi)}\delta_{y_j}$ is $q_{k,k+1}=\frac{K_{j_{k+1}}}{N}$ and the rate of leaving the state $Y_k=\frac{1}{N}\sum_{j\in J^k(\xi)}\delta_{y_j}$ is $q_k=\frac{1}{2}\sum_{\substack{j,l\in J^k(\xi)\\ \text{with } j\neq l}}\frac{K(y_j,y_l)}{N}$.

Hence,

$$
\begin{aligned}
&\mathbb{P}_y\left(S_i \le t < T_i^{[n]}, \xi_t^i \in d\xi\right)\\
&= \mathbb{P}_y\left(J_1 \in ds_1, \dots, J_{n-1} \in ds_{n-1}, Y_1 = \frac{1}{N}\sum_{j\in J^0(\xi)} \delta_{y_j}, \dots, Y_{n-1} = \frac{1}{N}\sum_{j\in J^{n-1}(\xi)} \delta_{y_j}\right)\\
&= q_{0,1}\dots q_{n-2,n-1}\exp\left(-q_1 s_1\right)\dots\exp\left(-q_{n-2}(s_{n-1}-s_{n-2})\right) ds_1\dots ds_{n-1}.
\end{aligned}
$$

Replacing q_k and $q_{k,k+1}$ by their respective values we obtain that

$$
\begin{aligned}
\mathbb{P}_y\left(S_i \le t < T_i^{[n]}, \xi_t^i \in d\xi\right) &= \left(\prod_{k=1}^{n-1}\frac{K_{j_k}}{N}\right)\exp\left(-\frac{1}{2}\sum_{\substack{j,l\in J^1(\xi)\\ \text{with } j\neq l}}\frac{K(y_j,y_l)}{N}s_1\right)\times\dots\\
&\qquad \dots\times\exp\left(-\frac{1}{2}\sum_{\substack{j,l\in J^{n-1}(\xi)\\ \text{with } j\neq l}}\frac{K(y_j,y_l)}{N}(s_{n-1}-s_{n-2})\right)\nu_i^y(d\xi)\\
&= \frac{K_\xi}{N^{n-1}}\exp\left(-\int_0^t \frac{K_s(\xi)}{N}ds\right)\nu_i^y(d\xi)
\end{aligned}
$$

as required.

□

Now let us prove Lemma 2.2. Before proving it, let us introduce some basic notations. For $\omega = \big(\omega_j : j \in \mathbb{T}_+^\star[N]\big) \in \Omega$, we will write $\omega^{[n]} = \big(\omega_j : j \in \mathbb{T}_+^\star[n]\big)$ and $\omega^J = \big(\omega_j : j \in \mathbb{T}_+^\star J\big)$.

Proof of Lemma 2.2: Fix $\xi \in A_i^y(0,t)$. We can find $\omega_0^{[n]} = (\omega_j^0 : j \in \mathbb{T}_+^\star[n])$ representing the configuration ξ. To ξ we can uniquely associate $\{(s_1,j_1),\dots,(s_{n-1},j_{n-1})\}$ where $s_1 < \dots < s_{n-1}$ are the coagulation times and $j_1,\dots,j_{n-1}$ are the labeled trees (subtrees of i) formed at $s_1,\dots,s_{n-1}$ respectively. Denote by $J^k(\xi)$ the set of particles from ξ present on $[s_k,s_{k+1})$ for $k\in\{0,\dots,n-1\}$ with the convention $s_0=0$ and $s_n=t$. For each $k\in\{1,\dots,n-1\}$, we write $j_k=\{j_k(1),j_k(2)\}$ with $j_k(1),j_k(2)\in J^{k-1}(\xi)$. Observe that $J^k(\xi)=\{j_k\}\cup J^{k-1}(\xi)\backslash\{j_k(1),j_k(2)\}$. Also, for convenience, for each $k\in\{1,\dots,n-1\}$ we will write

$$K_{j_k} = K(y_{j_k(1)}, y_{j_k(2)}).$$

Now, fix $\omega_0^J = (\omega_i^0 : i \in \mathbb{T}_+^\star J)$. There exist $r_1(\omega_0^J) < \dots < r_{n'-1}(\omega_0^J)$ such that $\mu_r^{N,J}(\omega_0^J)$ constant on $[r_k(\omega_0^J), r_{k+1}(\omega_0^J))$ for $k=0,\dots,n'-1$ with the convention $r_0(\omega_0^J)=0$ and $r_{n'}(\omega_0^J)=t$. For each $k\in\{0,\dots,n'-1\}$ denote by $J_k(\omega_0^J)$ the set of particles from $\mu_r^{N,J}(\omega_0^J)$ present on $[r_k(\omega_0^J), r_{k+1}(\omega_0^J))$. Define

$$\Omega(\omega_0^J,\xi) = \{\omega\in\Omega : \omega^{[n]} = \omega_0^{[n]}, \omega^J = \omega_0^J\}.$$

To obtain our result, conditional on $\{S_i \le t < T_i^{[n]}\}$ it is enough to compute :

$$\mathbb{P}_{y,\mu_0^{N,J}}\left(\{\omega\in\Omega(\omega_0^J,\xi) : t < T_i(\omega)\}\right) = \mathbb{P}_{y,\mu_0^{N,J}}\left(t < T_i(\omega_0^J,\omega_0^{[n]})\right) \tag{16}$$

for short where $T_i(\omega_0^J,\omega_0^{[n]})$ is understood as a random variable on $\Omega(\omega_0^J,\xi)$. Since we are conditioning on $\{S_i \le t < T_i^{[n]}\}$, on each interval $[s_k,s_{k+1})$, the particles present at time s_k in ξ that is $J^k(\xi)$, will not interfere between each other until time s_{k+1}. Hence, conditional on $\{S_i \le t < T_i^{[n]}\}$, the quantity (16) means that on each interval $[s_k,s_{k+1})$, each particle in $J^k(\xi)$ will not interfere with the configuration $\big(\mu_r^{N,J}(\omega_0^J)\big)_{r\ge s_k}$ until s_{k+1}. This is equivalent to consider on $[s_k,s_{k+1})$, for each $i_0\in J^k(\xi)$, starting from

$$\mu_0^{J_k(s_k)\cup\{i_0\}} = \sum_{j\in J_k(s_k)\cup\{i_0\}} \delta_{y_j},$$

that i_0 will survive until s_{k+1} where $J_k(s_k)$ is the set of particles present in $\mu_{s_k}^{N,J}(\omega_0^J)$ (this set is one of the $J_p(\omega_0^J)$ for $p\in\{0,\dots,n'-1\}$). If $T_{i_0}^{J_k(s_k)\cup\{i_0\}}$ denotes the time of death of the particle i_0 when starting from the set of particles $J_k(s_k)\cup\{i_0\}$ (this death time is defined in 1.4) we want $T_{i_0}^{J_k(s_k)\cup\{i_0\}}(\omega) > s_{k+1}$ for all $\omega\in\Omega(\omega_0^J,\xi)$. It is exactly the result we have obtained in Theorem 2.1. For short we will write, $T_{i_0}^{J_k(s_k)\cup\{i_0\}} = T_{i_0}^{s_k}$. Thus, setting

$$\begin{aligned}
A_1(\omega_0^J,\xi) &= \{\omega\in\Omega(\omega_0^J,\xi) : \forall i\in J^0(\xi), T_i^{s_1}(\omega) > s_1\}\\
&\vdots\\
A_{n-1}(\omega_0^J,\xi) &= \{\omega\in\Omega(\omega_0^J,\xi) : \forall i\in J^{n-2}(\xi), T_i^{s_{n-1}}(\omega) > s_{n-1}\}\\
A_n(\omega_0^J,\xi) &= \{\omega\in\Omega(\omega_0^J,\xi) : T_{j_{n-1}}^t(\omega) > t\},
\end{aligned}$$

we can write, conditional on $\{S_i \le t < T_i^{[n]}\}$,

$$\mathbb{P}_{y,\mu_0^{N,J}}\left(t < T_i(\omega_0^J,\omega^{[n]})\right) = \mathbb{P}_{y,\mu_0^{N,J}}\left(A_1(\omega_0^J,\xi)\cap\ldots\cap A_n(\omega_0^J,\xi)\right).$$

The process we consider is a continuous-time Markov process whose state space is the space of measures on the set $(0,\infty)$. Hence, fixing the coagulation times $s_1<\cdots<s_{n-1}$ we can consider the process independently on each interval $[s_k,s_{k+1})$ starting respectively from $(y^k,\mu_{s_k}^{N,J}(\omega_0^J))$ where $y^0=(y_1,\dots,y_n)$ and for $k\in\{0,\dots,n-1\}$, $y^k=(y_j, j\in J^k(\xi))$. Thus,

$$\begin{aligned}
\mathbb{P}_{y,\mu_0^{N,J}}\left(t < T_i(\omega_0^J,\omega_0^{[n]})\right) = \mathbb{P}_{y,\mu_0^{N,J}(\omega_0^J)}\left(A_1(\omega_0^J,\xi)\right)\times\mathbb{P}_{y^1,\mu_{s_1}^{N,J}(\omega_0^J)}\left(A_2(\omega_0^J,\xi)\right)\times\ldots\\
\ldots\times\mathbb{P}_{y^{n-1},\mu_{s_{n-1}}^{N,J}(\omega_0^J)}\left(A_n(\omega_0^J,\xi)\right).
\end{aligned}$$

First, let us compute

$$\mathbb{P}_{y,\mu_0^{N,J}(\omega_0^J)}\left(A_1(\omega_0^J,\xi)\right) = \mathbb{P}\left(\{\omega\in\Omega(\omega_0^J,\xi) : \forall i\in J^0(\xi), T_i^J(\omega) > s_1\}\right).$$

Take $i_0\in J^0(\xi)$. For $j\in J_k(\omega_0^J)$ for $k\in\{1,\dots,n'\}$ either $S_{\{i_0,j\}} = V_{\{i_0,j\}}$ if j is a simple particle, or there exist $p\in\{0,\dots,k\}$ such that $S_j(\omega_0^J) = r_p(\omega_0^J)$ if j is a composed particle. In this case $S_{\{i_0,j\}} = r_p(\omega_0^J) + V_{\{i_0,j\}}$. Hence, when $i_0\in\{1,\dots,n\}$, $T_{i_0}^J(\omega_0^J,\omega_0^{[n]})$ only depends on $(V_{\{i_0,j\}} : j\in J_k(\omega_0^J), k=0,\dots,n'-1)$. Hence, $\left(T_j^J(\omega_0^J,\omega_0^{[n]}) : j\in J^0(\xi)\right)$ are independent random variables. Thus,

$$\mathbb{P}_{y,\mu_0^{N,J}(\omega_0^J)}\left(A_1(\omega_0^J,\xi)\right) = \prod_{j\in J^0(\xi)} \mathbb{P}_{y,\mu_0^{N,J}(\omega_0^J)}\left(\{\omega\in\Omega(\omega_0^J,\xi) : T_j^J(\omega) > s_1\}\right).$$

For $j \in J^0(\xi)$, by Theorem 2.1,

$$\mathbb{P}_{y,\mu_0^{N,J}(\omega_0^J)}(\{\omega \in \Omega(\omega_0^J,\xi) : T_j^J(\omega) > s_1\}) = \exp\left(-\int_0^{s_1}\int_E K(y_j,y')\mu_r^{N,J}(\omega_0^J)(dy')dr\right).$$

Hence,

$$\mathbb{P}_{y,\mu_0^{N,J}(\omega_0^J)}\left(A_1(\omega_0^J,\xi)\right) = \exp\left(-\int_0^{s_1}\int_E \sum_{j\in J^0(\xi)} K(y_j,y')\mu_r^{N,J}(\omega_0^J)(dy')dr\right).$$

Now let us compute $\mathbb{P}_{y^k,\mu_{s_k}^{N,J}(\omega_0^J)}\left(A_k(\omega_0^J,\xi)\right)$ for $k \in \{2,\dots,n-1\}$. We are starting at time s_{k-1} with the set of particles $J^{k-1}(\xi)$. By the memoryless property for exponentially distributed random variables, it is equivalent to start off at time 0 with the set $J^{k-1}(\xi)$ and to go to $s_k - s_{k-1}$. Hence, the computation of $\mathbb{P}_{y^k,\mu_{s_k}^{N,J}(\omega_0^J)}(A_k(\omega_0^J,\xi))$ is the same than for $\mathbb{P}_{y,\mu_0^{N,J}(\omega_0^J)}\left(A_1(\omega_0^J,\xi)\right)$ replacing $J^0(\xi)$ by $J^{k-1}(\xi)$, j_1 by j_k, and s_1 by $s_k - s_{k-1}$. Thus,

$$\mathbb{P}_{y^k,\mu_{0s_k}^{N,J}(\omega_0^J)}(A_k(\omega_0^J,\xi)) = \exp\left(-\int_{s_{k-1}}^{s_k}\int_E \sum_{j\in J^{k-1}(\xi)} K(y_j,y')\mu_r^{N,J}(\omega_0^J)(dy')dr\right).$$

Thus,

$$\begin{aligned}
&\mathbb{P}_{y,\mu_0^{N,J}}\left(\{\omega \in \Omega(\omega_0^J,\xi) : t < T_i(\omega)\}\right) \\
&\quad = \prod_{k=1}^{n} \exp\left(-\int_{s_{k-1}}^{s_k}\int_E \sum_{j\in J^{k-1}(\xi)} K(y_j,y')\mu_r^{N,J}(\omega_0^J)(dy')dr\right) \\
&\quad = \exp\left(\sum_{k=1}^{n} -\int_{s_{k-1}}^{s_k}\int_E \sum_{j\in J^{k-1}(\xi)} K(y_j,y')\mu_r^{N,J}(\omega_0^J)(dy')dr\right) \\
&\quad = \exp\left(-\int_{\Delta(\xi)}\int_E K(y_r,y')\mu_{\Pi(r)}^{N,J}(dy')dr\right).
\end{aligned}$$

Hence,

$$\mathbb{P}_{y,\mu_0^{N,J}}\left(t < T_i | \mathcal{F}_J \vee \mathcal{F}_{[n]}\right) = \exp\left(-\int_{\Delta(\xi_t^i)}\int_E K(y_r,y')\mu_{\Pi_{\xi_t^i}(r)}^{N,J}(dy')dr\right) \text{ a.s}$$

as required.

□

Hence,

$$\begin{aligned}
&\mathbb{E}_{y,\mu_0^{N,J}}\left(f_i\left(\xi_t^i\right)1_{\{S_i\le t<T_i\}} \mid \mathcal{F}_J\right)N^{n-1} \\
&= \int_{A_i^y(0,t)} f_i(\xi)\,K_\xi \exp\left(-\int_{\Delta(\xi)}\int_E K(y_r,y')\mu_{\Pi_\xi(r)}^{N,J}(dy')dr\right)\exp\left(-\int_0^t \frac{K_s(\xi)}{N}ds\right)\nu_i^y(d\xi) \\
&= \int_{A_i^y(0,t)} f_i(\xi)\,K_\xi h_N\left(\left(\mu_r^{N,J}\right)_{r<t},\xi\right)\nu_i^y(d\xi)
\end{aligned}$$

where

$$h_N\left(\left(\mu_r^{N,J}\right)_{r<t},\xi\right)=\exp\left(-\int_{\Delta(\xi)}\int_E K(y_r,y')\mu^{N,J}_{\Pi_\xi(r)}(dy')dr\right)\exp\left(-\int_0^t\frac{K_s(\xi)}{N}ds\right)$$

and where for $y\in(0,\infty)^n$, the Lebesgue measure ν_i^y is defined in the Appendix 5.2.

2.4.3 Convergence for teh conditional expectation

Fix $y=(y_1,\dots,y_n)\in(0,\infty)^n$. This is the vector of masses associated to the particles $\{1,\dots,n\}$. Let $\tilde y$ be the tree vector of masses associated to $i\in\mathbb{T}_n^\star[n]$. Define

$$\begin{aligned}P_tf_i(y)=P_tf_i(\tilde y)&=\int_{A_i^y(0,t)}f_i\left(\xi\right)K_\xi\exp\left(-\int_{\Delta(\xi)}\int_E K(y_r,y')\mu_{\Pi_\xi(r)}(dy')dr\right)\nu_i^y(d\xi)\\&=\int_{A_i^y(0,t)}f_i\left(\xi\right)K_\xi h\left(\left(\mu_r\right)_{r<t},\xi\right)\nu_i^y(d\xi)\end{aligned}$$

where

$$h\left(\left(\mu_r\right)_{r<t},\xi\right)=\exp\left(-\int_{\Delta(\xi)}\int_E K(y_r,y')\mu_{\Pi_\xi(r)}(dy')dr\right).$$

Set

$$P_t^Nf_i(y)=P_t^Nf_i(\tilde y)=\mathbb{E}_{y,\mu_0^{N,J}}\left(f_i\left(\xi_t^i\right)1_{\{S_i\le t<T_i\}}\mid\mathcal{F}_J\right)N^{n-1}.$$

Theorem 2.4.

$P_t^Nf_i(y)\to P_tf_i(y)$ in probability as $N\to\infty$.

Proof of Theorem 2.4 : For a given $\xi\in A_i^y(0,t)$, we can write

$$\begin{aligned}h_N\left(\left(\mu_r^{N,J}\right)_{r<t},\xi\right)=&\prod_{k=0}^{n-1}\exp\left(-\frac12\sum_{\substack{j,l\in J^k(\xi)\\\text{with } j\neq l}}\frac{K(y_j,y_l)}{N}(s_{k+1}-s_k)\right)\\&\prod_{k=0}^{n-1}\exp\left(-\int_{s_k}^{s_{k+1}}\int_E\sum_{j\in J^k(\xi)}K(y_j,y')\mu_r^{N,J}(dy')dr\right)\end{aligned}$$

where $s_1<\cdots<s_{n-1}$ are the coagulation times associated to ξ and $j_1,\dots,j_{n-1}$ are its labeled subtrees (subtrees of i) formed at these respective times. $J^k(\xi)$ represents the set of tree particles (subtrees of i) from ξ present on $[s_k,s_{k+1})$ for $k=0,\dots,n-1$ with the convention $s_0=0$ and $s_n=t$. As $N\to\infty$, for each $k\in\{1,\dots,n-1\}$,

$$\prod_{k=0}^{n-1}\exp\left(-\frac12\sum_{\substack{j,l\in J^k(\xi)\\\text{with } j\neq l}}\frac{K(y_j,y_l)}{N}(s_{k+1}-s_k)\right)\to 1.$$

Now we want to show that

$$\prod_{k=0}^{n-1} \exp\left(-\int_{s_k}^{s_{k+1}} \int_E \sum_{j\in J^k(\xi)} K(y_j,y')\mu_r^{N,J}(dy')dr\right)$$
$$\to \prod_{k=0}^{n-1} \exp\left(-\int_{s_k}^{s_{k+1}} \int_E \sum_{j\in J^k(\xi)} K(y_j,y')\mu_r(dy')dr\right)$$

as $N\to\infty$ in probability. It is sufficient to show that for each $k\in\{1,\dots,n-1\}$ and $j\in J^k(\xi)$,

$$\exp\left(-\int_{s_k}^{s_{k+1}}\int_E K(y_j,y')\mu_r^{N,J}(dy')dr\right)\to \exp\left(-\int_{s_k}^{s_{k+1}}\int_E K(y_j,y')\mu_r(dy')dr\right)$$

as $N\to\infty$ in probability. For $k\in\{1,\dots,n-1\}$ and $j\in J^k(\xi)$ consider the random variable

$$X_N^{k,j} = \int_{s_k}^{s_{k+1}} K(y_j,y')\mu_r^{N,J}(dy')dr - \int_{s_k}^{s_{k+1}} K(y_j,y')\mu_r(dy')dr.$$

On $[s_k,s_{k+1})$,

$$\mu_r^N = \mu_r^{N,J} + \frac{1}{N}\sum_{l\in J_k(\xi)} \delta_{y_l}.$$

Hence ,

$$X_N^{k,j} = \int_{s_k}^{s_{k+1}} K(y_j,y')\mu_r^N(dy')dr$$
$$-\int_{s_k}^{s_{k+1}} K(y_j,y')\mu_r(dy')dr - (s_{k+1}-s_k)\sum_{l\in J_k(\xi)}\frac{K(y_j,y_l)}{N}.$$

Now, we know that

$$K(y_j,y) = \tilde{K}(y_j,y)\varphi(y_j)\varphi(y).$$

For $y\in(0,\infty)$ define $f_{y_j}:(0,\infty)\to(0,\infty)$ such that

$$f_{y_j}(y) = \tilde{K}(y_j,y)\varphi(y_j).$$

By hypothesis, this function is continuous and bounded. Hence, we can write

$$X_N^{k,j} = \int_{s_k}^{s_{k+1}} \langle f_{y_j}\varphi,\mu_r^N\rangle dr - \int_{s_k}^{s_{k+1}} \langle f_{y_j}\varphi,\mu_r\rangle dr - (s_{k+1}-s_k)\sum_{l\in J_k(\xi)}\frac{K(y_j,y_l)}{N}.$$

Hence ,

$$\left|X_N^{k,j}\right| \le \left|(s_{k+1}-s_k)\sum_{l\in J_k(\xi)}\frac{K(y_j,y_l)}{N}\right| + (s_{k+1}-s_k)\sup_{r\in[s_k,s_{k+1})}\left|\langle f_{y_j}\varphi,\mu_r^N\rangle - \langle f_{y_j}\varphi,\mu_r\rangle\right|.$$

The first term converges to 0. By (6), the second converges to 0 in probability as $N \to \infty$. Hence $X_N^{k,j} \to 0$ in probability as $N \to \infty$. Hence, for all $\xi \in A_i^y(0,t)$,

$$h_N\Big(\left(\mu_r^{N,J}\right)_{r<t}, \xi\Big) \to h\Big((\mu_r)_{r<t}, \xi\Big)$$

in probability as $N \to \infty$.

Now, on $A_i^y(0,t)$ K_ξ is bounded, say by some $C > 0$. Thus,

$$\left|P_t^N f_i(y) - P_t f_i(y)\right| \le C\|f_i\|_\infty \int_{A_i^{\tilde{y}}(0,t)} \left| h_N\left(\left(\mu_r^{N,J}\right)_{r<t}, \xi\right) - h\Big((\mu_r)_{r<t}, \xi\Big)\right| d\xi$$

Moreover, for all $\xi \in A_i^y(0,t)$,

$$\left| h_N\left(\left(\mu_r^{N,J}\right)_{r<t}, \xi\right)\right| \le 1.$$

Hence by the Bounded Convergence Theorem,

$$\mathbb{E}\left(\int_{A_i^y(0,t)} \left| h_N\left(\left(\mu_r^{N,J}\right)_{r\le t}, \xi\right) - h\Big((\mu_r)_{r\le t}, \xi\Big)\right| d\xi \right) \to 0$$

as $N \to \infty$. Thus,

$$\mathbb{E}\left(\left|P_t^N f_i(y) - P_t f_i(y)\right|\right) \to 0$$

as $N \to \infty$ and a fortiori

$$P_t^N f_i(y) \to P_t f_i(y)$$

as $N \to \infty$ in probability as required.

□

3 Some convergence results

The aim of this section is to prove (9, that is, for all $\tau \in \mathbb{T}$, for all $f \in C_b\left(A_\tau(0,t)\right)$,

$$\langle f, \tilde{\mu}_t^N\rangle \to \langle f, \tilde{\mu}_t\rangle$$

as $N \to \infty$ in probability.

3.1 Convergence of the expectation

We recall that we are working with the set of initial particles $[N] = \{1, \dots, N\}$ with associated masses $y_1, \dots, y_N > 0$. Let $\tau \in \mathbb{T}$ with n leaves, that is $n(\tau) = n$. Take $i \in \mathbb{T}_n^\star[n]$ with type τ. Associate the map $I : [N]_{[n]}^\star \to \mathbb{T}_n^\star[N]$ as defined in 2.1. For $(i_1, \dots, i_n) \in [N]_{[n]}^\star$ with associated masses $y_{i_1}, \dots, y_{i_n} > 0$ we will write $y_{(i_1,\dots,i_n)}$ for the vector of masses and $\tilde{y}_{(i_1,\dots,i_n)}$ for the vector tree of masses associated to $I(i_1, \dots, i_n)$. Let us fix $f \in C_b\left(A_\tau(0,t)\right)$. We aim to prove the following result.

Proposition 3.1.

$$\mathbb{E}\left(\langle f, \tilde{\mu}_t^N \rangle\right) \to \langle f, \tilde{\mu}_t \rangle$$

as $N \to \infty$*.*

Proof of Proposition 3.1 : For $M > 0$, define $\Psi_M : (0,\infty) \to (0,\infty)$ to be

$$\Psi_M(x) = \begin{cases} 1 & \text{if } x \le M \\ -x + (M+1) & \text{if } M \le x \le M+1 \\ 0 & \text{if } x \ge M+1 \end{cases}$$

We can write,

$$\langle f, \tilde{\mu}_t^N \rangle = \langle f \Psi_M(m), \tilde{\mu}_t^N \rangle + \langle f(1 - \Psi_M(m)), \tilde{\mu}_t^N \rangle$$

where $m : A(0,t) \to (0,\infty)$ is the mass function defined in subsection 1.3. Hence,

$$\mathbb{E}\left(\langle f, \tilde{\mu}_t^N \rangle\right) = \mathbb{E}\left(\langle f \Psi_M(m), \tilde{\mu}_t^N \rangle\right) + \mathbb{E}\left(\langle f(1 - \Psi_M(m)), \tilde{\mu}_t^N \rangle\right). \tag{17}$$

We use the following lemma that we shall prove below.

Lemma 3.2.

$$\mathbb{E}\left(\langle f \Psi_M(m), \tilde{\mu}_t^N \rangle\right) \to \langle f \Psi_M(m), \tilde{\mu}_t \rangle$$

as $N \to \infty$*.*

Thus, taking the limsup and liminf over N in the expression (17) and applying Lemma 3.2, we obtain,

$$\limsup_N \mathbb{E}\left(\langle f, \tilde{\mu}_t^N \rangle\right) = \langle f \Psi_M(m), \tilde{\mu}_t \rangle + \limsup_N \mathbb{E}\left(\langle f(1 - \Psi_M(m)), \tilde{\mu}_t^N \rangle\right) \tag{18}$$

and

$$\liminf_N \mathbb{E}\left(\langle f, \tilde{\mu}_t^N \rangle\right) = \langle f \Psi_M(m), \tilde{\mu}_t \rangle + \liminf_N \mathbb{E}\left(\langle f(1 - \Psi_M(m)), \tilde{\mu}_t^N \rangle\right). \tag{19}$$

Since $\mu_t^N = \tilde{\mu}_t^N \circ m^{-1}$ and $\varphi \ge 1$, we can write

$$\begin{aligned} \left|\langle f(1 - \Psi_M(m)), \tilde{\mu}_t^N \rangle\right| &= \left|\langle f(1 - \Psi_M), \mu_t^N \rangle\right| \\ &\le \|f\|_\infty \left|\langle \varphi(1 - \Psi_M), \mu_t^N \rangle\right| \end{aligned}$$

Hence,

$$\mathbb{E}\left(\left|\langle f(1 - \Psi_M(m)), \tilde{\mu}_t^N \rangle\right|\right) \le \|f\|_\infty \mathbb{E}\left(\left|\langle \varphi(1 - \Psi_M), \mu_t^N \rangle\right|\right). \tag{20}$$

Now, $\left|\langle \varphi(1 - \Psi_M), \mu_t^N \rangle\right|$ is bounded for all N. Indeed, since φ is sublinear, for all t, we have

$$\langle \varphi, \mu_t^N \rangle \le \langle \varphi, \mu_0^N \rangle.$$

Hence,

$$\langle \varphi(1 - \Psi_M), \mu_t^N \rangle \le \langle \varphi, \mu_0^N \rangle.$$

Now fix $\epsilon > 0$. Since $\langle \varphi, \mu_0^N \rangle \to \langle \varphi, \mu_0 \rangle$ as $N \to \infty$, we can find $N_0 > 0$ such that for all $N \ge N_0$,

$$\langle \varphi, \mu_0^N \rangle \le \langle \varphi, \mu_0 \rangle + \epsilon.$$

Let $M_0 = \max_{N \leq N_0} \langle \varphi, \mu_0^N \rangle$ and set $C_0 = \max(M_0, \langle \varphi, \mu_0 \rangle + \epsilon)$. For all N we have,

$$\langle \varphi, \mu_t^N \rangle \leq C_0$$

and a fortiori,

$$\langle \varphi(1 - \Psi_M), \mu_t^N \rangle \leq C_0$$

so it is bounded for all N. Moreover, it converges to $|\langle \varphi(1 - \Psi_M), \mu_t \rangle|$ in probability as $N \to \infty$. Thus, by the Bounded convergence Theorem,

$$\begin{aligned} \mathbb{E}\left(\left| \langle \varphi(1 - \Psi_M), \mu_t^N \rangle \right| \right) &\to \mathbb{E}\left(|\langle \varphi(1 - \Psi_M), \mu_t \rangle| \right) \\ &= |\langle \varphi(1 - \Psi_M), \mu_t \rangle| \end{aligned}$$

as $N \to \infty$. Taking the limsup over N in the expression (20) we obtain

$$\limsup_N \mathbb{E}\left(\left| \langle f(1 - \Psi_M(m)), \tilde{\mu}_t^N \rangle \right| \right) \leq \|f\|_\infty \left| \langle \varphi(1 - \Psi_M), \mu_t \rangle \right|.$$

As $M \to \infty$, $\varphi(1 - \Psi_M) \searrow 0$ and is positive. Hence, by the Monotone Convergence Theorem,

$$|\langle \varphi(1 - \Psi_M), \mu_t \rangle| \to 0 \text{ as } M \to \infty.$$

So

$$\lim_M \limsup_N \mathbb{E}\left(\left| \langle f(1 - \Psi_M(m)), \tilde{\mu}_t^N \rangle \right| \right) = \lim_M \liminf_N \mathbb{E}\left(\left| \langle f(1 - \Psi_M(m)), \tilde{\mu}_t^N \rangle \right| \right) = 0$$

Moreover,

$$|\langle f(1 - \Psi_M(m)), \tilde{\mu}_t \rangle| \leq \|f\|_\infty \left| \langle \varphi(1 - \Psi_M), \mu_t \rangle \right|$$

that is going to 0 as $M \to \infty$. Thus,

$$\lim_{M \to \infty} \langle f \Psi_M(m), \tilde{\mu}_t \rangle = \langle f, \tilde{\mu}_t \rangle.$$

Hence, taking the limit as $M \to \infty$ in the relations (18) and (19) we obtain

$$\limsup_N \mathbb{E}\left(\langle f, \tilde{\mu}_t^N \rangle \right) = \liminf_N \mathbb{E}\left(\langle f, \tilde{\mu}_t^N \rangle \right) = \langle f, \tilde{\mu}_t \rangle.$$

Thus,

$$\lim_N \mathbb{E}\left(\langle f, \tilde{\mu}_t^N \rangle \right) = \langle f, \tilde{\mu}_t \rangle$$

as required.

□

Proof of Lemma 3.2 : Set $f^M = f \Psi_M(m)$. We can write,

$$\mathbb{E}\left(\langle f^M, \tilde{\mu}_t^N \rangle \right) = \frac{2^{-q(\tau)}}{N} \sum_{(i_1, \ldots, i_n) \in [N]^\star_{[n]}} \mathbb{E}\left(f^M \left(\xi_t^{I(i_1, \ldots, i_n)} \right) 1_{\{S_{I(i_1, \ldots, i_n)} \leq t < T_{I(i_1, \ldots, i_n)}\}} \right)$$

where $I : [N]^\star_{[n]} \to \mathbb{T}^\star_n[N]$ is the map defined in subsection 2.1. For $(i_1, \dots, i_n) \in [N]^\star_{[n]}$, let $f^M_{I(i_1,\dots,i_n)}$ be the corresponding element for f^M in $C_b(A_{I(i_1,\dots,i_n)}(0,t))$ as defined in subsection 2.2. Then,

$$\begin{aligned}
&\mathbb{E}\left(\langle f^M, \tilde{\mu}^N_t \rangle\right) \\
&= \frac{1}{N} \sum_{(i_1,\dots,i_n)\in[N]^\star_{[n]}} \mathbb{E}\left(f^M_{I(i_1,\dots,i_n)}\left(\xi_t^{I(i_1,\dots,i_n)}\right) 1_{\{S_{I(i_1,\dots,i_n)} \le t < T_{I(i_1,\dots,i_n)}\}} \right) \\
&= \frac{1}{N^n} \sum_{(i_1,\dots,i_n)\in[N]^\star_{[n]}} \mathbb{E}\Bigg[\int_{A^{y(i_1,\dots,i_n)}_{I(i_1,\dots,i_n)}(0,t)} f^M_{I(i_1,\dots,i_n)}(\xi)\, K_\xi \\
&\exp\left(- \int_{\Delta(\xi)} \int_E K(y_r, y') \mu^{N,[N]\setminus\{i_1,\dots,i_n\}}_{\Pi_\xi(r)}(dy')dr \right) \exp\left(- \int_0^t \frac{K_s(\xi)}{N} ds \right) \nu^{y(i_1,\dots,i_n)}_{I(i_1,\dots,i_n)}(d\xi) \Bigg]
\end{aligned}$$

using the expression of the conditional expectation obtained at the end of the subsection 2.4.2.

Step 1: We are going to give an alternative expression of $\mathbb{E}\left(\langle f^M, \tilde{\mu}^N_t \rangle\right)$ using a new measure. Define on $(0, \infty)^n$ the measure,

$$\mu^{n,N}_0 = \frac{1}{N^n} \sum_{(i_1,\dots,i_n)\in[N]^\star_{[n]}} \delta_{(y_{i_1},\dots,y_{i_n})}.$$

For $(i_1, \dots, i_n) \in [N]^\star_{[n]}$ define

$$\begin{aligned}
&G(y_{i_1}, \dots, y_{i_n}) \\
&= \mathbb{E}\Bigg[\int_{A^{y(i_1,\dots,i_n)}_{I(i_1,\dots,i_n)}(0,t)} f^M_{I(i_1,\dots,i_n)}(\xi)\, K_\xi \exp\left(- \int_0^t \frac{K_s(\xi)}{N} ds \right) \\
&\exp\left(- \int_{\Delta(\xi)} \int_E K(y_r, y') \mu^{N,[N]\setminus\{i_1,\dots,i_n\}}_{\Pi_\xi(r)}(dy')dr \right) \nu^{y(i_1,\dots,i_n)}_{I(i_1,\dots,i_n)}(d\xi) \Bigg].
\end{aligned}$$

In this expression, the first three terms in the integral (inside the expectation) only depends on $(y_{i_1}, \dots, y_{i_n})$, whereas the last term depend on $(y_1, \dots, y_N)$. Nevertheless we only need to know which particles are present in the first three terms to work out which particles are going to be in the last term. Moreover, if there exist $i_1, i'_1 \in [N]$ with $i_1 \neq i'_1$ but $y_{i_1} = y_{i'_1}$ then we have

$$G(y_{i'_1}, y_{i_2} \dots, y_{i_n}) = G(y_{i_1}, y_{i_2} \dots, y_{i_n}).$$

Thus we can write,

$$\begin{aligned}
\mathbb{E}\left(\langle f^M, \tilde{\mu}^N_t \rangle\right) &= \frac{1}{N^n} \sum_{(i_1,\dots,i_n)\in[N]^\star_{[n]}} G(y_{i_1}, \dots, y_{i_n}) \\
&= \int_{(0,\infty)^n} G(x_1, \dots, x_n) \mu^{n,N}_0(dx_1 \dots dx_n).
\end{aligned}$$

Step 2: We are going to show that we can write

$$\mathbb{E}\left(\langle f^M, \tilde{\mu}_t^N\rangle\right) = \int_{(0,\infty)^n} G(x_1,\ldots,x_n)\mu_0^N(dx_1)\ldots\mu_0^N(dx_n) + \circ(\frac{1}{N}).$$

Let us consider

$$\left(\mu_0^N\right)^{\otimes n} = \left(\frac{1}{N}\sum_{i=1}^N \delta_{y_i}\right)^n = \mu_0^{n,N} + C_0^{n,N}$$

where $C_0^{n,N}$ is a finite sum of terms of the form

$$\mu_0^{N,n_1,\ldots,n_k} = \frac{1}{N^n} \sum_{\substack{i_1,\ldots,i_k=1\\ \text{with } i_1,\ldots,i_k\\ \text{distinct}}}^N \delta_{(\underbrace{y_{i_1},\ldots,y_{i_1}}_{n_1\ times},\underbrace{y_{i_2},\ldots,y_{i_2}}_{n_2\ times},\ldots,\underbrace{y_{i_k},\ldots,y_{i_k}}_{n_k\ times})}$$

for $k > 1$ with $n_1 + n_2 + \ldots + n_k = n$ modulo some permutations of the masses. Take $k < n-1$ and consider,

$$a_N^k = \frac{1}{N^n} \sum_{\substack{i_1,\ldots,i_k=1\\ \text{with } i_1,\ldots,i_k\\ \text{distinct}}}^N \mathbb{E}\Bigg(f^M\left(\xi_t^{I(i_1,\ldots,i_1,\ldots,i_k,\ldots,i_k)}\right)$$

$$1_{\{S_{I(i_1,\ldots,i_1,\ldots,i_k,\ldots,i_k)}\leq t< T_{I(i_1,\ldots,i_1,\ldots,i_k,\ldots,i_k)}\}}\Bigg)N^{n-1}.$$

where $I(i_1,\ldots,i_1,\ldots,i_k,\ldots,i_k)$ is obtained from $I(i_1,\ldots,i_n)$ with $(i_1,\ldots,i_n) \in [N]^{\star}_{[n]}$ by substituting in the tree $i(i_1,\ldots,i_n)$, the particles $\{i_p : 1 \leq p \leq n_1\}$ by the particle i_1, the particles $\{i_p : n_1+1 \leq p \leq n_1+n_2\}$ by the particle $i_2\ldots$, and finally the particles $\{i_p : n-n_k+1 \leq p \leq n\}$ by the particle i_k. It is a tree where particles are not all distinct. We are going to prove that this quantity is $\circ(\frac{1}{N})$. By a similar argument to before we can write

$$a_N^k = \frac{1}{N^n} \sum_{\substack{i_1,\ldots,i_k=1\\ \text{with } i_1,\ldots,i_k\\ \text{distinct}}}^N g(y_{i_1},\ldots y_{i_k})$$

where $g : (0,\infty)^k \to \mathbb{R}$ is the expectation above.

Now define

$$\left(\mu_0^N\right)^{\otimes k} = \left(\frac{1}{N}\sum_{i=1}^N \delta_{y_i}\right)^k$$

and on $(0,\infty)^k$ the measure,

$$\mu_0^{k,N} = \frac{1}{N^k} \sum_{\substack{i_1,\ldots,i_k=1\\ \text{with } i_1,\ldots,i_k\\ \text{distinct}}}^N \delta_{(y_{i_1},\ldots,y_{i_k})}.$$

Then,

$$\mu_0^{k,N} \leq \left(\mu_0^N\right)^{\otimes k}.$$

We can identify $\mu_0^{N,n_1,\ldots,n_k}$ and $\mu_0^{k,N}$. Hence,

$$\begin{aligned}
a_N^k &= \frac{1}{N^{n-k}} \int_{(0,\infty)^k} g(x_1, \ldots x_k) \mu_0^{k,N}(dx_1 \ldots dx_k) \\
&\le \frac{1}{N^{n-k}} \int_{(0,\infty)^k} g(x_1, \ldots x_k) \mu_0^{N}(dx_1) \ldots \mu_0^{N}(dx_k)
\end{aligned}$$

For $i \in \mathbb{T}_n^\star[n]$, let $A_{i,M}(0,t) = \{\xi \in A_i(0,t) : m(\xi) \le M+1\}$. On $A_{i,M}(0,t)$, the map $A(0,t) \to (0,\infty) : \xi \to K_\xi$ is bounded by $C = \left(\sup_{y,y' \in [0,M]^2} K(y,y')\right)^{n-1}$ (which is attained as K is bounded on $[0,M]^2$ compact). Outside $A_{i,M}(0,t)$, $\Psi_M(m)$ is the zero-function and so is f_M. Now, for all $(i_1, \ldots, i_n) \in [N]^n$ non necessarily distinct,

$$g(y_{i_1}, \ldots, y_{i_k}) = \mathbb{E}\left(P_t^N f^M_{I(i_1,\ldots,i_n)}\left(\tilde{y}_{(i_1,\ldots,i_n)}\right)\right).$$

Now,

$$\begin{aligned}
&\left|P_t^N f^M_{I(i_1,\ldots,i_n)}\left(\tilde{y}_{(i_1,\ldots,i_n)}\right)\right| \\
&= \Big| \int_{A^{y_{(i_1,\ldots,i_n)}}_{I(i_1,\ldots,i_n)}(0,t)} f^M_{I(i_1,\ldots,i_n)}(\xi)\, K_\xi \exp\left(-\int_0^t \frac{K_s(\xi)}{N} ds\right) \\
&\quad \exp\left(-\int_{\Delta(\xi)} \int_E K(y_r, y') \mu^{N,J}_{\Pi_\xi(r)}(dy') dr\right) \nu^{y_{(i_1,\ldots,i_n)}}_{I(i_1,\ldots,i_n)}(d\xi) \Big| \\
&\le C\|f\|_\infty \int_{A^{y_{(i_1,\ldots,i_n)}}_{I(i_1,\ldots,i_n)}(0,t)} \nu^{y_{(i_1,\ldots,i_n)}}_{I(i_1,\ldots,i_n)}(d\xi) \\
&= C\|f\|_\infty \int_{(0,\infty)^n} \sum_{\substack{\text{Possible permutations} \\ \text{for } s_1,\ldots,s_{n-1}}} 1_{\{s_1 \le \cdots \le s_{n-1}\}} ds_1 \ldots ds_{n-1} \\
&\le C\|f\|_\infty t^{n-1}.
\end{aligned}$$

Hence,

$$\begin{aligned}
a_N^k &\le \frac{C\|f\|_\infty t^{n-1}}{N^{n-k}} \int_{(0,\infty)^k} \mu_0^N(dx_1) \ldots \mu_0^N(dx_k) \\
&\le \frac{C\|f\|_\infty t^{n-1}}{N^{n-k}} \int_{(0,\infty)^k} \varphi(x_1)\mu_0^N(dx_1) \ldots \varphi(x_n)\mu_0^N(dx_k) \\
&= \frac{C\|f\|_\infty t^{n-1}}{N^{n-k}} \langle \varphi, \mu_0^N \rangle^k.
\end{aligned}$$

as $\varphi \ge 1$. It is clear that this quantity is going to 0 as $N \to \infty$ since $k < n-1$ and $\langle \varphi, \mu_0^N \rangle \to \langle \varphi, \mu_0 \rangle < \infty$ as $N \to \infty$. Hence,

$$\mathbb{E}\left(\langle f_M, \tilde{\mu}_t^N \rangle\right) = \int_{(0,\infty)^n} G(x_1, \ldots, x_n) \mu_0^N(dx_1) \ldots \mu_0^N(dx_n) + \circ(\frac{1}{N}).$$

as required.

Step 3: Let $i_0 \in \mathbb{T}_n^\star[n]$ with type τ and without associated masses. For each $(x_1, \dots, x_n) \in (0,\infty)^n$, we can associate to i_0 the vector of masses $(x_1, \dots, x_n)$. Hence we can write

$$G(x_1, \dots, x_n) = \mathbb{E}\left(P_t^N f_{i_0}^M\,(x_1, \dots, x_n)\right)$$

By Theorem 2.4, for all $x_1, \dots, x_n \in (0,\infty)$,

$$P_t^N f_{i_0}^M\,(x_1, \dots, x_n) \to P_t f_{i_0}^M\,(x_1, \dots, x_n)$$

in probability as $N \to \infty$. Also, $P_t^N f_{i_0}^M(x_1, \dots, x_n)$ is bounded by $C\|f\|_\infty t^{n-1}$. Hence, by the Bounded Convergence Theorem,

$$\mathbb{E}\left(\langle f_M, \tilde{\mu}_t^N\rangle\right) \to \int_{(0,\infty)^n} P_t f_{i_0}^M\,((x_1, \dots, x_n))\,\mu_0(dx_1)\dots\mu_0(dx_n)$$

as $N \to \infty$.

Now, we need to prove that

$$\int_{(0,\infty)^n} P_t f_{i_0}^M\,((x_1, \dots, x_n))\,\mu_0(dx_1)\dots\mu_0(dx_n) = \langle f^M, \tilde{\mu}_t\rangle.$$

But,

$$\begin{aligned}
&\int_{(0,\infty)^n} P_t f_{i_0}^M\,(x_1, \dots, x_n)\,\mu_0(dx_1)\dots\mu_0(dx_n) \\
&\quad = \int_{(0,\infty)^n} \int_{A_{i_0}^{(x_1,\dots,x_n)}(0,t)} K_\xi f_{i_0}^M\,(\xi)\,\nu_{i_0}^{(x_1,\dots,x_n)}(d\xi)\mu_0(dx_1)\dots\mu_0(dx_n) \\
&\qquad \exp\left(-\int_{\Delta(\xi)}\int_E K(y_r, y')\mu_r(dy')dr\right) \\
&\quad = \int_{A_{i_0}(0,t)} f_{i_0}^M\,(\xi)\,\tilde{\mu}'_t(d\xi) = \int_{A_\tau(0,t)} f^M\,(\xi)\,\tilde{\mu}_t(d\xi) = \langle f^M, \tilde{\mu}_t\rangle
\end{aligned}$$

using Appendix 5.3. Hence,

$$\mathbb{E}(\langle f^M, \tilde{\mu}_t^N\rangle) \to \langle f^M, \tilde{\mu}_t\rangle$$

as $N \to \infty$ as required.

□

3.2 Convergence of the expectation of the square

We are still working with $[N] = \{1, \dots, N\}$ with associated masses $y_1, \dots, y_N > 0$. Take $\tau \in \mathbb{T}$ with n leaves that is $n(\tau) = n$ for some $n > 0$. Fix $f \in C_b(A_\tau(0,t))$. The aim in this section is to prove the following result.

Proposition 3.3.

$$\mathbb{E}\left(\langle f, \tilde{\mu}_t^N\rangle^2\right) \to \langle f, \tilde{\mu}_t\rangle^2$$

as $N \to \infty$.

In this aim, we are going to proceed similarly to sections 2 and 3. First we are going to express $\mathbb{E}\left(\langle f, \tilde{\mu}_t^N\rangle^2\right)$ as the expected value of a finite sum of conditional expectations, then we will compute these conditional expectations and give some properties about their convergence. Finally we will prove Proposition 3.3.

3.2.1 An expression for the expectation

Take $i \in \mathbb{T}^\star_n[n]$ with type τ and associate the function $I : [N]^\star_{[n]} \to \mathbb{T}^\star_n[N]$ defined in 2.1. The aim is to compute

$$\mathbb{E}\left(\langle f, \tilde{\mu}^N_t \rangle^2\right).$$

We can write,

$$\begin{aligned}
\mathbb{E}\left(\langle f, \tilde{\mu}^N_t \rangle^2\right) &= \mathbb{E}\left(\left(\frac{2^{-q(\tau)}}{N} \sum_{\sigma \in S_{n,N}} f\left(\xi^{\sigma(i)}_t\right) 1_{\{S_{\sigma(i)} \le t < T_{\sigma(i)}\}}\right)^2\right) \\
&= \frac{2^{-2q(\tau)}}{N^2} \sum_{\sigma \in S_{n,N}} \mathbb{E}\Bigg(f^2\left(\xi^{\sigma(i)}_t\right) 1_{\{S_{\sigma(i)} \le t < T_{\sigma(i)}\}}\Bigg) \\
&+ \frac{2^{-2q(\tau)}}{N^2} \sum_{\substack{\sigma,\sigma' \in S_{n,N} \\ \text{with } Im(\sigma) \cap Im(\sigma') = \emptyset}} \mathbb{E}\Bigg(f\left(\xi^{\sigma(i)}_t\right) f\left(\xi^{\sigma'(i)}_t\right) 1_{\{S_{\sigma(i)} \le t < T_{\sigma(i)}\}} 1_{\{S_{\sigma'(i)} \le t < T_{\sigma'(i)}\}}\Bigg).
\end{aligned}$$

where $Im(\sigma) = \{\sigma(j) : j \in [N]\}$. Observe that in the formula above we omitted terms of the form

$$\mathbb{E}\Bigg(f\left(\xi^{\sigma(i)}_t\right) f\left(\xi^{\sigma'(i)}_t\right) 1_{\{S_{\sigma(i)} \le t < T_{\sigma(i)}\}} 1_{\{S_{\sigma'(i)} \le t < T_{\sigma'(i)}\}}\Bigg)$$

for $\sigma, \sigma' \in S_{n,N}$ with $Im(\sigma) \cap Im(\sigma') \neq \emptyset$. Indeed, in this case, by the way we constructed the $\{S_j : j \in \mathbb{T}^\star_+[N]\}$, the events $\{S_{\sigma(i)} \le t < T_{\sigma(i)}\}$ and $\{S_{\sigma'(i)} \le t < T_{\sigma'(i)}\}$ are disjoint (because by construction the trees have all distinct leaves) and so the quantity above is equal to 0. Let us look at the first term in the expression of $\mathbb{E}\left(\langle f, \tilde{\mu}^N_t \rangle^2\right)$. We have

$$\frac{2^{-2q(\tau)}}{N^2} \sum_{\sigma \in S_{n,N}} \mathbb{E}\Bigg(f^2\left(\xi^{\sigma(i)}_t\right) 1_{\{S_{\sigma(i)} \le t < T_{\sigma(i)}\}}\Bigg) = \frac{2^{-q(\tau)}}{N} \mathbb{E}\left(\langle f^2, \tilde{\mu}^N_t \rangle\right).$$

From subsection 3.1, as $N \to \infty$,

$$\mathbb{E}\left(\langle f^2, \tilde{\mu}^N_t \rangle\right) \to \langle f^2, \tilde{\mu}_t \rangle < \infty.$$

Hence as $N \to \infty$,

$$\frac{2^{-2q(\tau)}}{N^2} \sum_{\sigma \in S_{n,N}} \mathbb{E}\Bigg(f^2\left(\xi^{\sigma(i)}_t\right) 1_{\{S_{\sigma(i)} \le t < T_{\sigma(i)}\}}\Bigg) \to 0.$$

For $\sigma, \sigma' \in S_{n,N}$ with $Im(\sigma) \cap Im(\sigma') = \emptyset$, set

$$J(\sigma(i), \sigma'(i)) = [N] \setminus \{\lambda(\sigma(i)), \lambda(\sigma'(i))\}.$$

Hence,

$$\begin{aligned}
\mathbb{E}\left(\langle f, \tilde{\mu}^N_t \rangle^2\right) = \mathbb{E}\Bigg(\frac{2^{-2q(\tau)}}{N^2} \sum_{\substack{\sigma,\sigma' \in S_{n,N} \\ \text{with } Im(\sigma) \cap Im(\sigma') = \emptyset}} \mathbb{E}\Bigg(f\left(\xi^{\sigma(i)}_t\right) f\left(\xi^{\sigma'(i)}_t\right) \\
1_{\{S_{\sigma(i)} \le t < T_{\sigma(i)}\}} 1_{\{S_{\sigma'(i)} \le t < T_{\sigma'(i)}\}} | \mathcal{F}^{J(\sigma(i), \sigma'(i))}\Bigg)\Bigg) + \circ(\frac{1}{N})
\end{aligned}$$

Thus, we need to compute the following conditional expectation

$$\mathbb{E}\left(f\left(\xi_t^{\sigma(i)}\right) f\left(\xi_t^{\sigma'(i)}\right) 1_{\{S_{\sigma(i)}\le t<T_{\sigma(i)}\}} 1_{\{S_{\sigma'(i)}\le t<T_{\sigma'(i)}\}} | \mathcal{F}^{J(\sigma(i),\sigma'(i))}\right)$$

for $\sigma, \sigma' \in S_{n,N}$ with $Im(\sigma)\cap Im(\sigma')=\emptyset$.

3.2.2 An expression for the conditional expectation

In order to compute $\mathbb{E}\left(\langle f, \tilde{\mu}_t^N\rangle^2\right)$, without loss of generality, by subsection 3.2.1, we need to compute

$$\mathbb{E}_{y^1,y^2,\mu_0^{N,J}}\left(f\left(\xi_t^{i_1}\right) f\left(\xi_t^{i_2}\right) 1_{\{S_{i_1}\le t<T_{i_1}\}} 1_{\{S_{i_2}\le t<T_{i_2}\}} | \mathcal{F}_J\right)$$

for $i_1 \in \mathbb{T}_n^\star[n], i_2 \in \mathbb{T}_n^\star[2n]\backslash[n]$ both with type τ. The quantities y^1 and y^2 represent the respective vector of masses (as defined in 2.2) for i_1 and i_2, $J=\{2n+1,\dots,N\}$ and $\mathcal{F}_J = \sigma\left(U_j : j\in\mathbb{T}_+^\star J\right)$. Define $f_{i_1} = f\circ g_{i_1}$ and $f_{i_2} = f\circ g_{i_2}$ where g_{i_1} and g_{i_2} are the map on forgetting labels defined in subsection 2.2. Then $f_{i_1}\in C_b\left(A_{i_1}(0,t)\right)$ and $f_{i_2}\in C_b\left(A_{i_2}(0,t)\right)$.

Step 1 : A symmetry argument

By a similar symmetry argument that the one we used in subsection 2.2, we can write

$$\begin{aligned}&\mathbb{E}_{y^1,y^2,\mu_0^{N,J}}\left(f\left(\xi_t^{i_1}\right) f\left(\xi_t^{i_2}\right) 1_{\{S_{i_1}\le t<T_{i_1}\}} 1_{\{S_{i_2}\le t<T_{i_2}\}} | \mathcal{F}_J\right)\\ &\quad = \mathbb{E}_{y^1,y^2,\mu_0^{N,J}}\left(f_{i_1}\left(\xi_t^{i_1}\right) f_{i_2}\left(\xi_t^{i_2}\right) 1_{\{S_{i_1}\le t<T_{i_1}\}} 1_{\{S_{i_2}\le t<T_{i_2}\}} | \mathcal{F}_J\right).\end{aligned}$$

Thus it is enough to compute

$$\mathbb{E}_{y^1,y^2,\mu_0^{N,J}}\left(f_{i_1}\left(\xi_t^{i_1}\right) f_{i_2}\left(\xi_t^{i_2}\right) 1_{\{S_{i_1}\le t<T_{i_1}\}} 1_{\{S_{i_2}\le t<T_{i_2}\}} | \mathcal{F}_J\right).$$

Step 2 : Simplifying the computation

Set $\mathcal{F}_{[n]} = \sigma\left(U_j : j\in\mathbb{T}_+^\star[n]\right)$ and $\mathcal{F}_{[2n]\backslash[n]} = \sigma\left(U_j : j\in\mathbb{T}_+^\star[2n]\backslash[n]\right)$ and look at

$$\mathbb{E}_{y^1,y^2,\mu_0^{N,J}}\left(f_{i_1}\left(\xi_t^{i_1}\right) f_{i_2}\left(\xi_t^{i_2}\right) 1_{\{S_{i_1}\le t<T_{i_1}\}} 1_{\{S_{i_2}\le t<T_{i_2}\}} | \mathcal{F}_J \vee \mathcal{F}_{[n]} \vee \mathcal{F}_{[2n]\backslash[n]}\right). \tag{21}$$

Observe that $f_{i_1}\left(\xi_t^{i_1}\right) 1_{\{S_{i_1}\le t<T_{i_1}^{[n]}\}}$ is measurable with respect to $\mathcal{F}_{[n]}$ and that $f_{i_2}\left(\xi_t^{i_2}\right) 1_{\{S_{i_2}\le t<T_{i_2}^{[2n]\backslash[n]}\}}$ is measurable with respect to $\mathcal{F}_{[2n]\backslash[n]}$. Moreover, $\{t<T_{i_1}\}\subset\{t<T_{i_1}^{[n]}\}$ and $\{t<T_{i_2}\}\subset\{t<T_{i_2}^{[2n]\backslash[n]}\}$. Thus, the quantity (21) is equal to

$$\begin{aligned}&f_{i_1}\left(\xi_t^{i_1}\right) f_{i_2}\left(\xi_t^{i_2}\right) 1_{\{S_{i_1}\le t<T_{i_1}^{[n]}\}} 1_{\{S_{i_2}\le t<T_{i_2}^{[2n]\backslash[n]}\}}\\ &\qquad \mathbb{E}_{y^1,y^2,\mu_0^{N,J}}\left(1_{\{t<T_{i_1}\}} 1_{\{t<T_{i_2}\}} | \mathcal{F}_J \vee \mathcal{F}_{[n]} \vee \mathcal{F}_{[2n]\backslash[n]}\right).\end{aligned}$$

Hence we can write,

$$
\begin{aligned}
&\mathbb{E}_{y^1,y^2,\mu_0^{N,J}}\left(f_{i_1}\left(\xi_t^{i_1}\right)f_{i_2}\left(\xi_t^{i_2}\right)1_{\{S_{i_1}\le t<T_{i_1}\}}1_{\{S_{i_2}\le t<T_{i_2}\}}|\mathcal{F}_J\right)\\
&\quad=\int_{A_{i_1}^{y^1}(0,t)\times A_{i_1}^{y^2}(0,t)} f_{i_1}\left(\xi_1\right)f_{i_2}\left(\xi_2\right)\\
&\qquad\mathbb{P}_{y^1,y^2,\mu_0^{N,J}}\Big(\{t<T_{i_1}\}\cap\{t<T_{i_2}\}|\mathcal{F}_J\vee\mathcal{F}_{[n]}\vee\mathcal{F}_{[2n]\setminus[n]}\Big)\left(\xi_1,\xi_2\right)\\
&\qquad\mathbb{P}_{y^1,y^2}\Big(S_{i_1}\le t<T_{i_1}^{[n]},S_{i_2}\le t<T_{i_2}^{[2n]\setminus[n]},\xi_t^{i_1}\in d\xi_1,\xi_t^{i_2}\in d\xi_2\Big).
\end{aligned}
$$

Lemma 3.4. *1. Conditional on the event* $\{S_{i_1}\le t<T_{i_1}^{[n]}\}\cap\{S_{i_2}\le t<T_{i_2}^{[2n]\setminus[n]}\}$,

$$
\begin{aligned}
&\mathbb{P}_{y^1,y^2,\mu_0^{N,J}}\Big(\{t<T_{i_1}\}\cap\{t<T_{i_2}\}|\mathcal{F}_J\vee\mathcal{F}_{[n]}\vee\mathcal{F}_{[2n]\setminus[n]}\Big)\\
&\quad=\exp\left(-\frac{1}{2}\int_0^t\frac{K_s(\xi_t^{i_1},\xi_t^{i_2})}{N}ds\right)\\
&\quad\exp\left(-\int_{\Delta(\xi_t^{i_1})}\int_E K(y_r,y')\mu^{N,J}_{\Pi_{\xi_t^{i_1}}(r)}(dy')dr\right)\\
&\quad\exp\left(-\int_{\Delta(\xi_t^{i_2})}\int_E K(y_r,y')\mu^{N,J}_{\Pi_{\xi_t^{i_2}}(r)}(dy')dr\right)
\end{aligned}
$$

where

$$
K_s(\xi_t^{i_1},\xi_t^{i_2})=-\frac{1}{2}\sum_{\substack{r\in\Pi_{\xi_1}^{-1}(s)\\ r'\in\Pi_{\xi_2}^{-1}(s)}}K(y_r,y_{r'}).
$$

2.

$$
\begin{aligned}
&\mathbb{P}_{y^1,y^2}\Big(S_{i_1}\le t<T_{i_1}^{[n]},S_{i_2}\le t<T_{i_2}^{[2n]\setminus[n]},\xi_t^{i_1}\in d\xi_1,\xi_t^{i_2}\in d\xi_2\Big)\\
&\quad=\frac{K_{\xi_1}K_{\xi_2}}{N^{2n-2}}\exp\left(-\int_0^t\frac{K_s(\xi_1)}{N}ds\right)\exp\left(-\int_0^t\frac{K_s(\xi_2)}{N}ds\right)\nu_{i_1}^{y^1}(d\xi_1)\nu_{i_2}^{y^2}(d\xi_2).
\end{aligned}
$$

In the lemma above, the quantity (1) is similar to the one in Lemma 2.2 except that here, it means that for $\xi_1\in A_{i_1}^{y^1}(0,t),\xi_2\in A_{i_2}^{y^2}(0,t)$ and $\omega^J=(\omega_i^J:i\in\mathbb{T}_+^\star J)$, the configurations ξ_1 and ξ_2 do not interfere with each other and are not killed by $\left(\mu_r^{N,J}\right)_{r\ge 0}(\omega^J)$. Hence, mimicking the proof of Lemma 2.2, we obtain the formula of the lemma. The second part of the proof comes from applying the Lemma 2.3 along with an argument of independence.

Hence,

$$\begin{aligned}
&\mathbb{E}_{y^1,y^2,\mu_0^{N,J}}\left(f_{i_1}\left(\xi_t^{i_1}\right)f_{i_2}\left(\xi_t^{i_2}\right)1_{\{S_{i_1}\le t<T_{i_1}\}}1_{\{S_{i_2}\le t<T_{i_2}\}}|\mathcal{F}_J\right)\\
&\quad=\int_{A_{i_1}^{y^1}(0,t)\times A_{i_2}^{y^2}(0,t)} f_{i_1}\left(\xi_1\right)f_{i_2}\left(\xi_2\right)\exp\left(-\frac{1}{2}\int_0^t\frac{K_s(\xi_1,\xi_2)}{N}ds\right)\\
&\qquad\frac{K_{\xi_1}K_{\xi_2}}{N^{2n-2}}\exp\left(-\int_0^t\frac{K_s(\xi_1)}{N}ds\right)\exp\left(-\int_0^t\frac{K_s(\xi_2)}{N}ds\right)\\
&\qquad\exp\left(-\int_{\Delta(\xi_1)}\int_E K(y_r,y')\mu^{N,J}_{\Pi_{\xi_1}(r)}(dy')dr\right)\\
&\qquad\exp\left(-\int_{\Delta(\xi_2)}\int_E K(y_r,y')\mu^{N,J}_{\Pi_{\xi_2}(r)}(dy')dr\right)\nu_{i_1}^{y^1}(d\xi_1)\nu_{i_2}^{y^2}(d\xi_2).
\end{aligned}$$

3.2.3 Convergence for the conditional expectation

We keep the same notations that in the previous section and we define,

$$P_t^N f_{i_1}(y^1)f_{i_2}(y^2)=\mathbb{E}_{y^1,y^2,\mu_0^{N,J}}\left(f_{i_1}\left(\xi_t^{i_1}\right)f_{i_2}\left(\xi_t^{i_2}\right)1_{\{S_{i_1}\le t<T_{i_1}\}}1_{\{S_{i_2}\le t<T_{i_2}\}}|\mathcal{F}_J\right)N^{2n-2}.$$

and

$$\begin{aligned}
&P_tf_{i_1}(y^1)f_{i_2}(y^2)\\
&=\int_{A_{i_1}^{y^1}(0,t)\times A_{i_2}^{y^2}(0,t)} K_{\xi_1}f_{i_1}(\xi_1)\exp\left(-\int_{\Delta(\xi_1)}\int_E K(y_r,y')\mu^{N,J}_{\Pi_{\xi_1}(r)}(dy')dr\right)\nu_{i_1}^{y^1}(d\xi_1)\\
&\qquad K_{\xi_2}f_{i_2}(\xi_2)\exp\left(-\int_{\Delta(\xi_2)}\int_E K(y_r,y')\mu^{N,J}_{\Pi_{\xi_2}(r)}(dy')dr\right)\nu_{i_2}^{y^2}(d\xi_2).
\end{aligned}$$

Proposition 3.5. *1. $P_t^N f_{i_1}(y^1)f_{i_2}(y^2)\to P_tf_{i_1}(y^1)f_{i_2}(y^2)$ as $N\to\infty$ in probability.*

2. $P_tf_{i_1}(y^1)f_{i_2}(y^2)=P_tf_{i_1}(y^1)P_tf_{i_2}(y^2)$.

We are not proving this proposition since the proof is similar to the proof of Theorem 2.4.

3.2.4 Proof of Proposition 3.3

We are going to sketch the proof of this proposition as it is very similar to the proof of Proposition 3.1.

Step 1: For $M>0$, define $\Psi_M:(0,\infty)\to(0,\infty)$ to be

$$\Psi_M(x)=\begin{cases}1 & \text{if } x\le M\\ -x+(M+1) & \text{if } M\le x\le M+1\\ 0 & \text{if } x\ge M+1\end{cases}$$

We can write,

$$\langle f,\tilde{\mu}_t^N\rangle=\langle f\Psi_M(m),\tilde{\mu}_t^N\rangle+\langle f(1-\Psi_M(m)),\tilde{\mu}_t^N\rangle.$$

Hence,

$$\langle f, \tilde{\mu}_t^N \rangle^2 = \langle f\Psi_M(m), \tilde{\mu}_t^N \rangle^2 + \langle f(1-\Psi_M(m)), \tilde{\mu}_t^N \rangle^2 + 2\langle f\Psi_M(m), \tilde{\mu}_t^N \rangle \langle f(1-\Psi_M(m)), \tilde{\mu}_t^N \rangle$$

and so

$$\begin{aligned}\mathbb{E}\left(\langle f, \tilde{\mu}_t^N \rangle^2\right) = \mathbb{E}\left(\langle f\Psi_M(m), \tilde{\mu}_t^N \rangle^2\right) &+ \mathbb{E}\left(\langle f(1-\Psi_M(m)), \tilde{\mu}_t^N \rangle^2\right) \\ &+ 2\mathbb{E}\left(\langle f\Psi_M(m), \tilde{\mu}_t^N \rangle \langle f(1-\Psi_M(m)), \tilde{\mu}_t^N \rangle\right).\end{aligned}$$

Step 2: We prove that $\mathbb{E}\left(\langle f\Psi_M(m), \tilde{\mu}_t^N \rangle^2\right) \to \langle f\Psi_M(m), \tilde{\mu}_t \rangle^2$ as $N \to \infty$. Let $i_1 \in \mathbb{T}_n^\star[n]$ and $i_2 \in \mathbb{T}_n^\star[2n]\backslash[n]$ both with type τ, and associate the maps $I_1 : [N]_{[n]}^\star \to \mathbb{T}_n^\star[N]$ and $I_2 : [N]_{[2n]\backslash[n]}^\star \to \mathbb{T}_n^\star[N]$ defined in subsection 2.1. For $M > 0$, let $f^M = f\Psi_M(m)$ where m is the mass function. Let $f_{i_1}^M$ and $f_{i_2}^M$ be the correspondent representant for f_M respectively in $C_b(A_{i_1}(0,t))$ and $C_b(A_{i_2}(0,t))$. Define on $(0,\infty)^{2n}$, the measure

$$\mu_0^{2n,N} = \frac{1}{N^{2n}} \sum_{\substack{i_1,\dots,i_n,j_1,\dots,j_n=1 \\ \text{with } i_1,\dots,i_n,j_1,\dots,j_n \\ \text{distinct}}}^{N} \delta_{(y_{i_1},\dots,y_{i_n},y_{j_1},\dots,y_{j_n})}.$$

We can write

$$\begin{aligned}&\mathbb{E}\left(\langle f^M, \tilde{\mu}_t^N \rangle^2\right) \\ &= \mathbb{E}\left(\int_{(0,\infty)^{2n}} P_t^N f_{i_1}^M\left((x_1,\dots,x_n)\right) f_{i_2}^M\left((x_{n+1},\dots,x_{2n})\right) \mu_0^{2n,N}(dx_1,\dots,dx_{2n})\right) \\ &\quad + \circ(\frac{1}{N})\end{aligned}$$

Hence, by mimicking Lemma 2.2 we can write,

$$\begin{aligned}&\mathbb{E}\left(\langle f^M, \tilde{\mu}_t^N \rangle^2\right) \\ &= \mathbb{E}\left(\int_{(0,\infty)^{2n}} P_t^N f_{i_1}^M\left((x_1,\dots,x_n)\right) f_{i_2}^M\left((x_{n+1},\dots,x_{2n})\right) \mu_0^N(dx_1),\dots,\mu_0^N(dx_{2n})\right) \\ &\quad + \circ(\frac{1}{N})\end{aligned}$$

By Theorem 3.4,

$$\begin{aligned}&P_t^N f_{i_1}^M\left((x_1,\dots,x_n)\right) f_{i_2}^M\left((x_{n+1},\dots,x_{2n})\right) \\ &\qquad \to P_t f_{i_1}^M\left((x_1,\dots,x_n)\right) f_{i_2}^M\left((x_{n+1},\dots,x_{2n})\right)\end{aligned}$$

as $N \to \infty$ and is bounded by $\|f_{i_1}^M\|_\infty \|f_{i_2}^M\|_\infty C^2 t^{2n-2}$ where C has been defined in 3.1. Hence by the Bounded Convergence Theorem,

$$\begin{aligned}\mathbb{E}\left(\langle f^M, \tilde{\mu}_t^N \rangle^2\right) \to \int_{(0,\infty)^{2n}} & P_t f_{i_1}^M\left((x_1,\dots,x_n)\right) f_{i_2}^M\left((x_{n+1},\dots,x_{2n})\right) \\ & \mu_0(dx_1)\dots\mu_0(dx_{2n})\end{aligned}$$

as $N \to \infty$. Now, by Theorem 3.4,

$$\begin{aligned} &P_t f_{i_1}^M ((x_1, \dots, x_n)) \, f_{i_2}^M ((x_{n+1}, \dots, x_{2n})) \\ &\qquad = P_t f_{i_1}^M ((x_1, \dots, x_n)) \, P_t f_{i_2}^M ((x_{n+1}, \dots, x_{2n})) \, . \end{aligned}$$

Hence, as $N \to \infty$,

$$\begin{aligned} \mathbb{E} \left(\langle f^M, \tilde{\mu}_t^N \rangle^2 \right) \to &\left(\int_{(0,\infty)^n} P_t f_{i_1}^M ((x_1, \dots, x_n)) \, \mu_0(dx_1) \dots \mu_0(dx_n) \right) \\ &\left(\int_{(0,\infty)^n} P_t f_{i_2}^M ((x_1, \dots, x_n)) \, \mu_0(dx_1) \dots \mu_0(dx_n) \right) \\ &= \langle f^M, \tilde{\mu}_t \rangle^2 . \end{aligned}$$

Step 3: We prove the Proposition 3.3. We have proved before that as $M \to \infty$, $\langle f^M, \tilde{\mu}_t \rangle \to \langle f, \tilde{\mu}_t \rangle$. Hence,

$$\langle f^M, \tilde{\mu}_t \rangle^2 \to \langle f, \tilde{\mu}_t \rangle^2$$

as $M \to \infty$. Also, similarly to the way that we proved that

$$\lim_M \limsup_N \mathbb{E} \left(\left| \langle f(1 - \Psi_M(m)), \tilde{\mu}_t^N \rangle \right| \right) = \lim_M \liminf_N \mathbb{E} \left(\left| \langle f(1 - \Psi_M(m)), \tilde{\mu}_t^N \rangle \right| \right) = 0$$

we prove that

$$\lim_M \limsup_N \mathbb{E} \left(\left| \langle f(1 - \Psi_M(m)), \tilde{\mu}_t^N \rangle \right|^2 \right) = \lim_M \liminf_N \mathbb{E} \left(\left| \langle f(1 - \Psi_M(m)), \tilde{\mu}_t^N \rangle \right|^2 \right) = 0.$$

Moreover,

$$\begin{aligned} &\mathbb{E} \left(\left| \langle f \Psi_M(m), \tilde{\mu}_t^N \rangle \langle f(1 - \Psi_M(m)), \tilde{\mu}_t^N \rangle \right| \right)^2 \\ &\qquad \leq \mathbb{E} \left(\langle f \Psi_M(m), \tilde{\mu}_t^N \rangle^2 \right) \mathbb{E} \left(\langle f(1 - \Psi_M(m)), \tilde{\mu}_t^N \rangle^2 \right) . \end{aligned}$$

Hence, we deduce that

$$\begin{aligned} &\lim_M \limsup_N \mathbb{E} \left(\left| \langle f \Psi_M(m), \tilde{\mu}_t^N \rangle \langle f(1 - \Psi_M(m)), \tilde{\mu}_t^N \rangle \right| \right)^2 \\ &\qquad = \lim_M \liminf_N \mathbb{E} \left(\left| \langle f \Psi_M(m), \tilde{\mu}_t^N \rangle \langle f(1 - \Psi_M(m)), \tilde{\mu}_t^N \rangle \right| \right)^2 = 0. \end{aligned}$$

Hence, we obtain

$$\limsup_N \mathbb{E} \left(\langle f, \tilde{\mu}_t^N \rangle^2 \right) = \liminf_N \mathbb{E} \left(\langle f, \tilde{\mu}_t^N \rangle^2 \right) = \langle f, \tilde{\mu}_t \rangle^2 .$$

Thus,

$$\lim_N \mathbb{E} \left(\langle f, \tilde{\mu}_t^N \rangle^2 \right) = \langle f, \tilde{\mu}_t \rangle^2$$

as required.

3.3 Conclusion

For all $\tau \in \mathbb{T}$ and for all $f \in C_b(A_\tau(0,t))$ we have proved that

$$\mathbb{E}\left(\langle f, \tilde{\mu}_t^N \rangle\right) \to \langle f, \tilde{\mu}_t \rangle$$

and

$$\mathbb{E}\left(\langle f, \tilde{\mu}_t^N \rangle^2\right) \to \langle f, \tilde{\mu}_t \rangle^2$$

as $N \to \infty$. So we deduce by the remark we did at the beginning of the paper, that for all $f \in C_b(A_\tau(0,t))$,

$$\langle f, \tilde{\mu}_t^N \rangle \to \langle f, \tilde{\mu}_t \rangle$$

as $N \to \infty$ in probability. Now let us prove our main result.

4 Proof of Theorem 1.1

Our aim in this section is to prove our main result stated in Theorem 1.1, that is

$$\tilde{\mu}_t^N \to \tilde{\mu}_t$$

as $N \to \infty$ weakly in probability. In Section 3, we have proved that for $\tau \in \mathbb{T}$ and for all $f \in C_b(A_\tau(0,t))$,

$$\langle f, \tilde{\mu}_t^N \rangle \to \langle f, \tilde{\mu}_t \rangle$$

as $N \to \infty$ in probability. To prove Theorem 1.1, we need to prove that for all $f \in C_b(A(0,t))$,

$$\langle f, \tilde{\mu}_t^N \rangle \to \langle f, \tilde{\mu}_t \rangle$$

as $N \to \infty$ in probability.

4.1 A tightness argument

We are reviewing here a particular case of the work of [1], [2]. The usual Marcus-Lushnikov process gives at each time the distribution in masses of the particles present in the system but does not retain any notion of configuration of these particles. We are going to define a process $(\overline{X}_t^N)_{t\geq 0}$ (the Marcus Lushnikov on trees) on $\mathbb{T}(0,\infty)$, the space of trees on $(0,\infty)$. The space $\mathbb{T}(0,\infty)$ is given by

$$\mathbb{T}(0,\infty) = \bigcup_{\tau \in \mathbb{T}} \mathbb{T}_\tau(0,\infty)$$

where $\mathbb{T}_1(0,\infty) = (0,\infty)$ and for $\tau = \{\tau_1, \tau_2\} \in \mathbb{T}$,

$$\mathbb{T}_\tau(0,\infty) = \big\{ y = \{y_1, y_2\} : y_1 \in \mathbb{T}_{\tau_1}(0,\infty), y_2 \in \mathbb{T}_{\tau_2}(0,\infty) \big\}.$$

On $\mathbb{T}(0,\infty)$ define the mass function $m' : \mathbb{T}(0,\infty) \to (0,\infty)$. For $y \in \mathbb{T}_1(0,\infty) = (0,\infty)$, we set

$$m'(y) = y.$$

Recursively for $\tau = \{\tau_1, \tau_2\} \in \mathbb{T}$, for $y = \{y_1, y_2\} \in \mathbb{T}_\tau(0, \infty)$, we set

$$m'(y) = m'(y_1) + m'(y_2).$$

Define also the counting function $n' : \mathbb{T}(0, \infty) \to \mathbb{N}$ as follows. For $y \in \mathbb{T}_1(0, \infty) = (0, \infty)$ we set

$$n'(y) = 1.$$

Recursively for $\tau = \{\tau_1, \tau_2\} \in \mathbb{T}$, for $y = \{y_1, y_2\} \in \mathbb{T}_\tau(0, \infty)$, we set

$$n'(y) = n'(y_1) + n'(y_2).$$

Let us define $(\overline{X}_t)_{t \geq 0}$. Let $[N] = \{1, \dots, N\}$ and let $y_1, \dots, y_N > 0$ be the masses associated to each particle in $[N]$. Set

$$\overline{X}_0 = \sum_{i=1}^{N} \delta_{y_i}.$$

For each $i < j \in [N]$ take an independent random variable T_{ij} such that T_{ij} is exponential with parameter $K\left(m'(y_i), m'(y_j)\right)$, and define

$$T = \min_{i<j} T_{ij}.$$

Set $\overline{X}_t = \overline{X}_0$ for $t < T$ and

$$\overline{X}_T = \overline{X}_t - (\delta_{y_i} + \delta_{y_j} - \delta_{\{y_i, y_j\}})$$

if $T = T_{ij}$, then begin the construction afresh from $\overline{X}_T$. Let $(\overline{X}_t^N)_{t \geq 0}$ be Marcus-Lushnikov on $\mathbb{T}(0, \infty)$ with kernel $\frac{K}{N}$ starting from

$$\overline{X}_0^N = \sum_{i=1}^{N} \delta_{y_i}.$$

Set

$$\overline{\mu}_t^N = N^{-1} \overline{X}_t^N.$$

Let

$$y_t : A(0, t) \to \mathbb{T}(0, \infty)$$

be the map on forgetting times. Hence $\tilde{\mu}_t^N$ and $\overline{\mu}_t^N$ are related through the following equality

$$\overline{\mu}_t^N = \tilde{\mu}_t^N \circ y_t^{-1}. \tag{22}$$

Define

$$\tilde{\varphi} : \mathbb{T}(0, \infty) \to (0, \infty) \text{ by } \tilde{\varphi} = \varphi \circ m.$$

On $\mathbb{T}_1(0, \infty) = (0, \infty)$, $\tilde{\varphi} = \varphi$ and so

$$\langle \tilde{\varphi}^2, \mu_0 \rangle = \langle \varphi^2, \mu_0 \rangle < \infty$$

by (4). Then [1] tells us that we can find $n_0 > 0$ such that for all N,

$$\mathbb{P}\left(\overline{\mu}_t^N\left(\{y \in \mathbb{T}(0,\infty) : n'(y) \geq n_0\}\right) > C(\epsilon)\}\right) < \epsilon$$

where $C(\epsilon) > 0$. Now by (22)

$$\overline{\mu}_t^N\left(\{y \in \mathbb{T}(0,\infty) : n'(y) \geq n_0\}\right) = \tilde{\mu}_t^N\left(\{\xi \in A(0,t) : n'\left(y_t(\xi)\right) \geq n_0\}\right).$$

Hence,

$$\mathbb{P}\left(\tilde{\mu}_t^N\left(\{\xi \in A(0,t) : n\left(y_t(\xi)\right) \geq n_0\}\right) > \frac{\epsilon}{\|f\|_\infty}\right) < \epsilon. \tag{23}$$

4.2 Proof of Theorem 1.1

Take $f \in C_b\left(A(0,t)\right)$. For $n \in \mathbb{N}$ consider

$$A_n = \bigcup_{\substack{\tau\in\mathbb{T} \\ \text{with } n(\tau) \leq n}} A_\tau(0,t).$$

Since

$$A(0,t) = \bigcup_{\tau\in\mathbb{T}} A_\tau(0,t)$$

it is clear that

$$A(0,t) = \bigcup_{n\geq 1} A_n.$$

Moreover, $(A_n)_{n\geq 1}$ forms an increasing sequence. Set $f_n = f 1_{A_n}$. Then, $f_n \in C_b\left(A_n\right)$ and $f_n \to f$ as $n \to \infty$. We can write

$$f_n = \sum_{\substack{\tau\in\mathbb{T} \\ \text{with } n(\tau) \leq n}} f_n 1_{A_\tau(0,t)}.$$

Thus,

$$\langle f_n, \tilde{\mu}_t^N\rangle = \sum_{\substack{\tau\in\mathbb{T} \\ \text{with } n(\tau) \leq n}} \langle f_n 1_{A_\tau(0,t)}, \tilde{\mu}_t^N\rangle.$$

Now, since $f_n 1_{A_\tau(0,t)} \in C_b\left(A_\tau(0,t)\right)$, by subsection 3.3,

$$\langle f_n 1_{A_\tau(0,t)}, \tilde{\mu}_t^N\rangle \to \langle f_n 1_{A_\tau(0,t)}, \tilde{\mu}_t\rangle$$

as $N \to \infty$, in probability. Thus, since the sum over $\{\tau \in \mathbb{T} \text{ with } n(\tau) \leq n\}$ is finite, we obtain

$$\langle f_n, \tilde{\mu}_t^N\rangle \to \langle f_n, \tilde{\mu}_t\rangle \tag{24}$$

as $N \to \infty$, in probability. Now let us prove our result. Also, for all n, $f_n \to f$ as $n \to \infty$ and $|f_n| \leq |f|$. Hence, by the dominated convergence theorem,

$$\tilde{\mu}_t(f_n) \to \tilde{\mu}_t(f) \tag{25}$$

as $n \to \infty$.

Now let us fix $\epsilon > 0$. Consider,

$$\begin{aligned}|\langle f, \tilde{\mu}_t^N \rangle - \langle f, \tilde{\mu}_t \rangle| \leq & |\langle f, \tilde{\mu}_t^N \rangle - \langle f_n, \tilde{\mu}_t^N \rangle| + |\langle f_n, \tilde{\mu}_t^N \rangle - \langle f_n, \tilde{\mu}_t \rangle| \\ & + |\langle f_n, \tilde{\mu}_t \rangle - \langle f, \tilde{\mu}_t \rangle|\end{aligned}$$

By subsection 4.1, we know that we can find $n_0 > 0$, such that for all N,

$$\mathbb{P}\left(\tilde{\mu}_t^N\left(\{\xi \in A(0,t) : n'\left(y_t(\xi)\right) \geq n_0\}\right) > \frac{\epsilon}{\|f\|_\infty}\right) < \frac{\epsilon}{3}. \tag{26}$$

Now, $f - f_n = f 1_{A_n^c}$. Thus,

$$|\langle f, \tilde{\mu}_t^N \rangle - \langle f_n, \tilde{\mu}_t^N \rangle| = |\langle f - f_n, \tilde{\mu}_t^N \rangle| \leq \|f\|_\infty |\langle 1_{A_n^c}, \tilde{\mu}_t^N \rangle|$$

For all $n \geq n_0, A_n^c \subseteq A_{n_0}^c$. So using the relation (25) we obtain that

$$\begin{aligned}\mathbb{P}\left(|\langle f, \tilde{\mu}_t^N \rangle - \langle f_n, \tilde{\mu}_t^N \rangle| > \epsilon\right) &< \mathbb{P}\left(\|f\|_\infty |\langle 1_{A_n^c}, \tilde{\mu}_t^N \rangle| > \epsilon\right) \\ &< \frac{\epsilon}{3}\end{aligned}$$

for all $n \geq n_0$. Hence, for $n \geq n_0$, for all N,

$$\mathbb{P}\left(|\langle f, \tilde{\mu}_t^N \rangle - \langle f_n, \tilde{\mu}_t^N \rangle| > \epsilon\right) < \frac{\epsilon}{3}$$

For the third term, by 4.1, we can choose n_1 so that, for all $n \geq n_1$,

$$|\langle f_n, \tilde{\mu}_t \rangle - \langle f, \tilde{\mu}_t \rangle| < \frac{\epsilon}{3}$$

For the second term : Take $n \geq \max(n_0, n_1)$. By the relation (24), we can find N_0 so that for all $N \geq N_0$,

$$\mathbb{P}\left(|\langle f_n, \tilde{\mu}_t^N \rangle - \langle f_n, \tilde{\mu}_t \rangle| > \epsilon\right) < \frac{\epsilon}{3}$$

Hence, for $N > N_0$,

$$\mathbb{P}\left(|\langle f, \tilde{\mu}_t^N \rangle - \langle f, \tilde{\mu}_t \rangle| > \epsilon\right) < \epsilon$$

that is

$$\langle f, \tilde{\mu}_t^N \rangle \to \langle f, \tilde{\mu}_t \rangle$$

as $N \to \infty$ in probability. Hence,

$$\tilde{\mu}_t^N \to \tilde{\mu}_t$$

as $N \to \infty$ weakly in probability as required.

5 Appendix

5.1 A topology on $A(0,t)$

We equip $A(0,t)$ with a topology. Define

$$\hat{}: A(0,t) \to A(0,1)$$

as follows. For $\xi \in A_1(0,t)$ we set $\hat{\xi} = \xi$. Recursively, for $\tau = \{\tau_1, \tau_2\} \in \mathbb{T}, \xi = (s, \{\xi_1, \xi_2\}) \in A_\tau(0,t)$ with $\xi_1 \in A_{\tau_1}(0,s)$ and $\xi_2 \in A_{\tau_2}(0,s)$ we define $\hat{\xi} = (\frac{s}{t}, \{\hat{\xi_1}, \hat{\xi_2}\}) \in A_\tau(0,1)$ with $\hat{\xi_1} \in A_{\tau_1}(0,1)$ and $\hat{\xi_2} \in A_{\tau_2}(0,1)$.

We are going to construct a topology on $A(0,1)$ and then rescaling we will obtain a topology on $A(0,t)$. The set $\mathbb{T}$ is countable so we can give it a strict total order $<$. For $\tau = 1 \in \mathbb{T}, A_1(0,1) = (0,\infty)$. We equip $A_1(0,1)$ with the usual topology on $(0,\infty)$. For $\tau = \{\tau_1, \tau_2\} \in \mathbb{T}$ with $\tau_1 < \tau_2$, we can identify $A_\tau(0,1)$ to be

$$A_\tau(0,1) = (0,1) \times A_{\tau_1}(0,1) \times A_{\tau_2}(0,1)$$

Now, for $\tau = \{1\}$,

$$A_{\{1\}}(0,1) = (0,1) \times A_1(0,1) \times A_1(0,1)$$

We equip $(0,1)$ with the usual topology on $\mathbb{R}$. We have already given $A_1(0,1)$ a topology. Hence, we equip $A_{\{1\}}(0,1)$ with the product topology (Tychonoff topology). By induction, assume we have topologies on $A_{\tau_1}(0,1)$ and $A_{\tau_2}(0,1)$ for $\tau_1, \tau_2 \in \mathbb{T}$. Recursively for $\tau = \{\tau_1, \tau_2\} \in \mathbb{T}$ with $\tau_1 < \tau_2$, we equip

$$A_\tau(0,1) = (0,1) \times A_{\tau_1}(0,1) \times A_{\tau_2}(0,1)$$

with the product topology. Then ,

$$A(0,1) = \bigcup_{\tau \in \mathbb{T}} A_\tau(0,1)$$

is naturally equipped with a topology. Finally, rescaling, we obtain a topology on $A(0,t)$ for $t \geq 0$. Similarly, we can equip $\tilde{A}(0,t)$ with a topology.

5.2 A measure on $A_i^y(0,t)$

Fix $0 \leq t < T$. The set $\mathbb{T}_n^\star[n]$ is countable so we can give it a total order $<$. Take $i = \{i_1, i_2\} \in \mathbb{T}_n^\star[n]$ with $i_1 < i_2$. Let y, y^1 and y^2 by the vector of masses respectively associated to i, i_1 and i_2. Our aim is to define a measure on $A_i^y(0,t)$. We are first going to construct a measure on $A_i^y(0,1)$ and then by a similar rescalling to the one done in subsection 5.1 we will obtain a measure on $A_i^y(0,t)$. Equip $A_i(0,1)$ with its Borel-σ algebra $\mathcal{B}_i$. We can We can identify $A_i^y(0,1)$ to be

$$A_i^y(0,1) = (0,1) \times A_{i_1}^{y^1}(0,1) \times A_{i_2}^{y^2}(0,1)$$

as $i_1 < i_2$. For $i = 1$, for $\xi \in A_i^y(0,1)$, set

$$\nu_i^y(d\hat{\xi}) = \delta_y.$$

Recursively for $i = \{i_1, i_2\} \in \mathbb{T}_n^\star[n]$ with $i_1 < i_2$, with associated vector of masses y, y^1 and y^2, for $\hat{\xi} = (s, \hat{\xi_1}, \hat{\xi_2}) \in A_i(0,1)$, define

$$\nu_i^y(d\hat{\xi}) = \nu_{i_1}^{y^1}(d\hat{\xi_1}) \nu_{i_2}^{y^2}(d\hat{\xi_2}) ds.$$

This defines a measure on the product space

$$\mathcal{B}\left((0,1)\right) \otimes \mathcal{B}_{i_1} \otimes \mathcal{B}_{i_2}$$

where $\mathcal{B}_{i_1}$ and $\mathcal{B}_{i_2}$ are the respective Borel-σ algebra on $A_{i_1}^{y^1}(0,1)$ and $A_{i_2}^{y^2}(0,1)$. Then rescalling as in 5.1 we obtain a measure on $A_i^y(0,t)$.

5.3 A limit measure on $\tilde{A}(0,t)$

Let $E = (0,\infty)$. Fix $0 \leq t < T$. Fix $n > 0$. We define a limit measure on $\tilde{A}(0,t)$ as follows. For $\xi \in A_k(0,t)$ with $k \in \mathbb{N}$, set

$$\tilde{\mu}'_t(d\xi) = \exp\left(-\int_0^t \int_E K(y,y')\mu_r(dy')\mathrm{d}r\right)\mu_0(dy)$$

where $y = m(\xi)$ and $(\mu_r)_{r<T}$ is the strong deterministic solution to the generalized Smoluchowski equation (3). For $i = \{i_1, i_2\} \in \mathbb{T}^\star_n\mathbb{N}$ with type $\tau(i) = \tau = \{\tau_1, \tau_2\} \in \mathbb{T}$, for $\xi = (s, \{\xi_1, \xi_2\}) \in A_i(0,t)$, with $s < t$, define recursively

$$\tilde{\mu}'_t(d\xi) = \epsilon(\tau)K(m(\xi_1), m(\xi_2))\tilde{\mu}'_s(d\xi_1)\tilde{\mu}'_s(d\xi_2)\exp\left(-\int_s^t \int_E K(y,y')\mu_r(dy')\mathrm{d}r\right)ds \quad (27)$$

where $y = m(\xi)$ and $\epsilon(\tau) = 1$ if $\tau_1 \neq \tau_2$ and $\epsilon(\tau) = \frac{1}{2}$ if $\tau_1 = \tau_2$.

For $\xi \in A_k(0,t)$ with $k \in \mathbb{N}$, set

$$K_\xi = 1$$

For $i = \{i_1, i_2\} \in \mathbb{T}^\star_n\mathbb{N}$, $\xi = (s, \{\xi_1, \xi_2\}) \in A_i(0,t)$, define recursively

$$K_\xi = K(m(\xi_1), m(\xi_2))K_{\xi_1}K_{\xi_2}$$

Theorem 5.1. *Let $i = \{i_1, i_2\} \in \mathbb{T}^\star_n\mathbb{N}$ with type $\tau = \tau(i) \in \mathbb{T}$.*

1. *Then for $\xi = (s, \{\xi_1, \xi_2\}) \in A_i(0,t)$,*

   $$\tilde{\mu}'_t(d\xi) = 2^{-q(\tau)}K_\xi \exp\left(-\int_{\Delta(\xi)}\int_E K(y_r, y')\mu_{\Pi(r)}(dy')dr\right)\nu^y_i(d\xi)\mu_0(dy_1)\ldots\mu_0(dy_n)$$

   where $y = (y_1, \ldots, y_n)$ is the vector of masses associated to ξ and ν^y_i is the measure described in subsection 5.2.

2. *Take $f \in C_b(A_\tau(0,t))$. Let $f_i = f \circ g_i$ where $g_i : A_i(0,t) \to A_\tau(0,t)$ is the map on forgetting labels. Then,*

   $$\int_{A_\tau(0,t)} f(\xi)\tilde{\mu}_t(d\xi) = \int_{A_i(0,t)} f_i(\xi)\int_{A_\tau(0,t)} f(\xi)\tilde{\mu}_t(d\xi)$$

Proof of Theorem 5.1:

(a) Let us do it by induction. It is clearly true for $i \in \mathbb{T}^\star_1\mathbb{N}$. Fix $n > 0$. Suppose it is true for all $i \in \mathbb{T}^\star_k\mathbb{N}$ with $k \leq n-1$. Is is true for $k = n$? Take $i = \{i_1, i_2\} \in \mathbb{T}^\star_n\mathbb{N}$. Then, writing $n(i_1) = k$ and $n(i_2) = n-k$ we have

$$n(i_1) \leq n-1$$

and

$$n(i_2) \leq n-1$$

Take $\xi = (s, \{\xi_1, \xi_2\}) \in A_i(0,t)$. Without loss of generality, assume that i_1 is formed from the particle $1, \dots, k$ with associated masses $y^1 = (y_1, \dots, y_k)$ and that i_2 is formed from the particle $k+1, \dots, n$ with associated masses $y^2 = (y_{k+1}, \dots, y_n)$. Using the induction hypothesis, we have,

$$\begin{aligned}\tilde{\mu}'_s(d\xi_1) =& 2^{-q(\tau_1)} K_{\xi_1} \exp\left(- \int_{\Delta(\xi_1)} \int_E K(y_r, y') \mu_{\Pi(r)}(dy') dr \right) \\ & \nu_{i_1}^{y^1}(d\xi_1) \mu_0(dy_1) \dots \mu_0(dy_k)\end{aligned}$$

and

$$\begin{aligned}\tilde{\mu}'_s(d\xi_2) =& 2^{-q(\tau_2)} K_{\xi_2} \exp\left(- \int_{\Delta(\xi_2)} \int_E K(y_r, y') \mu_{\Pi(r)}(dy') dr \right) \\ & \nu_{i_2}^{y^2}(d\xi_2) \mu_0(dy_{k+1}) \dots \mu_0(dy_n)\end{aligned}$$

Using the relation (27), we have

$$\begin{aligned}\tilde{\mu}'_t(d\xi) =& \epsilon(\tau) 2^{-q(\tau_1)} 2^{-q(\tau_2)} K(m(\xi_1), m(\xi_2)) K_{\xi_1} K_{\xi_2} \nu_{i_1}^{y^1}(d\xi_1) \nu_{i_2}^{y^2}(d\xi_2) ds \mu_0(dy_1) \dots \mu_0(dy_n) \\ & \exp\left(- \int_{\Delta(\xi_1)} \int_E K(y_r, y') \mu_{\Pi(r)}(dy') dr \right) \exp\left(- \int_{\Delta(\xi_2)} \int_E K(y_r, y') \mu_{\Pi(r)}(dy') dr \right) \\ & \exp\left(- \int_s^t \int_E K(y, y') \mu_r(dy') \mathrm{d}r \right)\end{aligned}$$

Now, $K_\xi = K(m(\xi_1), m(\xi_2)) K_{\xi_1} K_{\xi_2}$, $\nu_i^y(d\xi) = \nu_{i_1}^{y^1}(d\xi_1) \nu_{i_2}^{y^2}(d\xi_2) ds$, $\Delta(\xi) = \Delta(\xi_1) \cup \Delta(\xi_2) \cup ((s,t] \times \xi)$ and $2^{-q(\tau)} = 2^{-q(\tau_1)} 2^{-q(\tau_2)} 2^{-\epsilon(\tau)}$. Hence,

$$\tilde{\mu}'_t(d\xi) = 2^{-q(\tau)} K_\xi \exp\left(- \int_{\Delta(\xi)} \int_E K(y_r, y') \mu_{\Pi(r)}(dy') dr \right) \nu_i^y(d\xi) \mu_0(dy_1) \dots \mu_0(dy_n)$$

as required.

□

5.4 Some simulations

Here are some simulations under Visual Basic of the Marcus-Lushnikov process on trees. The graphics below represent trees that have been simulated following the Marcus-Lushnikov process on trees with different kernel K and an initial number of particles N. In these simulations, all the initial particles have mass 1. These pictures show for each kernel the sort of trees limit we can expect to find in the limit measure.



Figure 1: $N = 128$, $K(x, y) = 1$



Figure 2: $N = 128$, $K(x, y) = xy$



Figure 3: $N = 128$, $K(x, y) = 1/(x + y + 1)$

References

[1] James Norris *Cluster Coagulation* Communications in Mathematical Physics, Vol. V209, No. 2. (1 February 2000), pp. 407-435

[2] James Norris *Smoluchowski's coagulation equation: uniqueness, non uniqueness and a hydrodynamic limit for the stochastic coalescent* Stochastic Process. Appl. 119 (2009), no. 1, 167–189. 60K35 (45K05 60H30 82C22)

[3] James Norris *Coagulation of Brownian particles*

[4] Nicolas Fournier, Philippe Laurencot *Marcus-Lushnikov processes, Smoluchowski's and Flory's models* Stochastic Process. Appl. 119 (2009), no. 1, 167–189.

[5] Nicolas Fournier and Jean-Sebastien Giet *Convergence of the Marcus-Lushnikov process* MSC 2000 : 45K05, 60H30.

[6] A.Marcus, Stochastic coalescence, Technometrics 10 (1968), pp. 133143.